A FINITE DIFFERENCE METHOD FOR A STEFAN PROBLEM

by

THOMAS CLAYTON SMITH, B.A., M.A.

A DISSERTATION

IN

MATHEMATICS

Submitted to the Graduate Faculty
of Texas Tech University in
Partial Fulfillment of
the Requirements for
the Degree of

DOCTOR OF PHILOSOPHY

Approved

May, 1978

AC
801
73
1978
No. 23
cop. 2

ACKNOWLEDGEMENTS

The author wishes to sincerely thank Dr. Harold D. Meyer for his guidance, his support, and his serving as the chairman of the committee. The author also wishes to thank, Dr. Wayne T. Ford, Dr. Richard E. Saeks, Dr. Charles N. Kellogg and Dr. Ronald Anderson for serving on his committee.

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	ii
LIST OF FIGURES	iv
INTRODUCTION	1
I. FORMULATION OF THE CONTINUOUS TIME STEFAN PROBLEM . . .	3
II. FORMULATION OF THE FINITE DIFFERENCE METHOD	5
III. SOME PRELIMINARY RESULTS	9
IV. CONVERGENCE	20
V. A NUMERICAL EXAMPLE	44
LIST OF REFERENCES	47

LIST OF FIGURES

Figure		Page
5.1	Comparison of Values for $U_{i,j}$	45
5.2	Comparison of Values for S_j	45

INTRODUCTION

Many physical problems involve initial boundary value problems for parabolic differential equations in which part of the boundary is not given a priori but is found as part of the solution. These problems have been considered under the name of "Stefan problems." Stefan problems occur in such physical processes as the melting of solids and the crystallizing of liquids.

We now give an example of such a problem from Friedman [4]. Consider a thin block of ice occupying an interval $0 \leq x < \infty$. We assume that the temperature of the ice is everywhere 0°C , and at the point $x = 0$ the temperature is kept at $T^\circ\text{C}$ where $T > 0$. Then the ice will begin to melt, and for every time $t > 0$ the water will occupy an interval $0 \leq x < s(t)$. Letting u denote the water temperature we have

$$(0.1) \quad \begin{cases} \alpha^2 u_{xx} - u_t = 0 & \text{for } 0 < x < s(t), t > 0, \\ u(0, t) = T & \text{for } t > 0, \\ u(s(t), t) = 0 & \text{for } t > 0, \end{cases}$$

where α is some constant not equal to zero, and $s(t)$ is the free boundary which is not given a priori. Furthermore, we have an additional condition on $x = s(t)$, namely, the law of conservation of energy. This condition is given by the equation

$$(0.2) \quad \frac{ds(t)}{dt} = -k u_x(s(t), t) \quad \text{for } t > 0,$$

where k is some positive constant.

The problem (0.1), (0.2) is called a Stefan problem. Existence and uniqueness for various one dimensional Stefan problems have been shown by several authors ([1], [4], [5], [8], [9]). Some multidimensional Stefan problems were considered by ([6], [11]).

The purpose of this paper is to approximate the solution of a Stefan problem using an implicit finite difference analog for the heat equation and an explicit finite difference analog for the differential equation describing the free boundary. Also, we shall show that the finite difference solution we obtain converges uniformly to the actual solution. Numerous numerical schemes for various Stefan problems have been successfully employed by several authors ([2], [3], [10], [12], [14], [16], [17]).

In Chapter I we formulate a continuous time Stefan problem, and in Chapter II we describe a finite difference scheme for approximating the solution of this problem. Chapter III contains some preliminary results. Chapter IV contains the main result which shows that the finite difference solution converges to the actual solution. Finally, in Chapter V we give a numerical example.

CHAPTER I

FORMULATION OF THE CONTINUOUS TIME STEFAN PROBLEM

The Stefan problem we are concerned with is the following: find two functions $s = s(t)$ and $u = u(x, t)$ such that

$$(1.1) \quad \left\{ \begin{array}{l} Lu \equiv u_{xx} - u_t = 0 \text{ in } 0 < x < s(t), 0 < t \leq T, \\ u(0, t) = f(t) \quad \text{for } 0 \leq t \leq T, \\ u(s(t), t) = 0 \quad \text{for } 0 \leq t \leq T, \\ u(x, 0) = \phi(x) \quad \text{for } 0 \leq x \leq b, \\ s(0) = b, \end{array} \right.$$

$$(1.2) \quad \frac{ds(t)}{dt} = -u_x(s(t), t) \quad \text{for } 0 < t \leq T,$$

where f , ϕ and b are given data, and T is a fixed but arbitrary positive number.

We make the following assumptions on f , ϕ and b : f and ϕ are nonnegative, b is positive, $f \in C^1$ for $0 \leq t \leq T$, $\phi \in C^2$ for $0 \leq x \leq b$ and there exists a positive constant c such that

$$(1.3) \quad 0 \leq \phi(x) \leq c(b-x) \quad \text{for } 0 \leq x \leq b.$$

In the above C^1 and C^2 are the usual spaces of functions possessing, respectively, continuous first and second derivatives in their given domains.

Let s be a given continuous function and assume that $s(t) > 0$ for $t \geq 0$. A solution of (1.1) is a function u , continuous for $0 \leq x \leq s(t)$, $0 \leq t \leq T$, such that in $0 < x < s(t)$, $0 < t \leq T$, u_{xx} and u_t are continuous and $Lu = 0$, and such that u satisfies the boundary conditions in (1.1).

A solution of (1.1), (1.2) is a pair of functions (s,u) such that $s(0) = b$, $s(t) > 0$, $s \in C^1$ for $0 < t \leq T$, and u satisfies (1.1); also such that $u_x(s(t),t)$ exists, is continuous, and satisfies (1.2).

Existence and uniqueness of the solution of (1.1), (1.2) have been established by Cannon and Hill [1]. Cannon and Hill have also shown that if s is Lipschitz continuous on $0 \leq t \leq T$, then (1.1), (1.2) is equivalent to (1.1), (1.4) where (1.4) is given by

$$(1.4) \quad s(t)^2 = b^2 + 2\int_0^t f(\tau)d\tau + 2\int_0^b \xi\phi(\xi)d\xi - 2\int_0^{s(t)} \xi u(\xi,t)d\xi.$$

CHAPTER II

FORMULATION OF THE FINITE DIFFERENCE SCHEME

In this chapter we formulate a finite difference scheme for approximating the solution of (1.1), (1.2). Let $0 = x_0 < x_1 < x_2 < \dots < x_M = b$ be a partition of the interval $[0, b]$ where $x_i = i\Delta x$, and let $0 = t_0 < t_1 < t_2 < \dots < t_N = T$ be a partition of the interval $[0, T]$ where $t_j = j\Delta t$. We let S_j denote the finite difference approximation of $s(t_j)$ and we let $U_{i,j}$ denote the finite difference approximation to $u_{i,j}$ where $u_{i,j} = u(x_i, t_j)$. The pair $(S_i, U_{i,j})$ is defined to be the solution of

$$(2.1) \left\{ \begin{array}{l} \frac{U_{i,j+1} - U_{i,j}}{\Delta t} = \frac{U_{i+1,j+1} - 2U_{i,j+1} + U_{i-1,j+1}}{(\Delta x)^2}, \\ \qquad \qquad \qquad i = 1, 1, 2, \dots, M-1, \\ \qquad \qquad \qquad j = 0, 1, 2, \dots, N-1, \\ \\ \frac{U_{M+i,j+1} - U_{M+i,j}}{\Delta t} = \frac{\frac{U_{M+i+1,j+1} - U_{M+i,j+1}}{\langle S_{i+1} \rangle - \langle S_i \rangle}}{\frac{1}{2}(\langle S_{i+1} \rangle - \langle S_{i-1} \rangle)}, \\ \\ \frac{U_{M+i,j+1} - U_{M+i-1,j+1}}{\langle S_i \rangle - \langle S_{i-1} \rangle} - \frac{1}{\frac{1}{2}(\langle S_{i+1} \rangle - \langle S_{i-1} \rangle)}, \\ \qquad \qquad \qquad j = J, J+1, J+2, \dots, N-1, \\ \qquad \qquad \qquad i = 0, 1, 2, \dots, L(j+1)-1, \end{array} \right.$$

$$(2.2) \quad \left\{ \begin{array}{l} \frac{S_1 - S_0}{\Delta t} = \frac{U_{M-1,0}}{\Delta x} = \frac{\phi(b-\Delta x)}{\Delta x}, \\ \frac{S_{j+1} - S_j}{\Delta t} = \frac{U_{M+L(j)-1,j}}{S_j - [S_{j-1}]}, \quad j = 1, 2, \dots, N-1, \end{array} \right.$$

$$(2.3) \quad \left\{ \begin{array}{l} U_{i,0} = \phi(i\Delta x), \quad i = 0, 1, 2, \dots, M, \\ U_{0,j} = f(j\Delta t), \quad j = 0, 1, 2, \dots, N, \\ U_{M+L(j),j} = 0, \quad j = 0, 1, 2, \dots, N, \end{array} \right.$$

where

$$J = \min\{k | S_k > S_0\} - 1,$$

$$L(j) = \begin{cases} 0 & \text{if } j = 0, \\ L(j-1) & \text{if } S_j = S_{j-1}, \\ L(j-1)+1 & \text{if } S_j \neq S_{j-1}, \end{cases}$$

$$\langle S_{-1} \rangle = x_{M-1},$$

$$\langle S_0 \rangle = S_0,$$

$$\langle S_i \rangle = \min\{S_k | S_k > \langle S_{i-1} \rangle\}, \quad i = 1, 2, \dots, L(N),$$

$$[S_j] = x_{M-1}, \quad j = 0, 1, \dots, J-1,$$

$$[S_j] = \max\{S_k | S_k < S_{j+1}\}, \quad j = J, J+1, \dots, N-1.$$

It is easy to see that this finite difference scheme has a unique solution. For each j , $j = 0, 1, 2, \dots, N-1$, (2.1) - (2.3) gives us a system of linear equations of the form $AU = V$, where

$$A = \begin{bmatrix} C(1,1) & C(1,2) & & 0 & . & . & . & . & 0 \\ C(2,1) & C(2,2) & C(2,3) & 0 & . & . & . & . & 0 \\ 0 & C(3,2) & C(3,3) & C(3,4) & & . & . & . & . \\ . & . & . & . & . & & & . & . \\ . & . & . & . & . & . & . & . & 0 \\ . & . & . & . & . & . & . & . & C(M+L(j+1)-2, M+L(j+1)-1) \\ 0 & . & . & . & 0 & C(M+L(j+1)-1, M+L(j+1)-2) & C(M+L(j+1)-1, M+L(j+1)-1) & . & . \end{bmatrix},$$

$$U = \begin{bmatrix} U_{1,j+1} \\ U_{2,j+1} \\ . \\ . \\ . \\ U_{M+L(j+1)-1,j+1} \end{bmatrix}, \quad V = \begin{bmatrix} U_{i,j} + rU_{0,j+1} \\ U_{2,j} \\ . \\ . \\ . \\ U_{M+L(j)-1,j} \\ 0 \end{bmatrix},$$

$$r = \frac{\Delta t}{(\Delta x)^2} ,$$

$$C(n,n) = 1+2r, \quad n = 1, 2, \dots, N-1,$$

$$C(n,n-1) = -r, \quad n = 2, 3, \dots, M-1,$$

$$C(n-1,n) = -r, \quad n = 2, 3, \dots, M,$$

$$C(M,M-1) = - \frac{2\Delta t}{\Delta x(\Delta x + \langle S_1 \rangle - S_0)} ,$$

$$C(M,M) = 1 + \frac{2\Delta t}{(\langle S_1 \rangle - S_0)(\Delta x + \langle S_1 \rangle - S_0)} + \frac{2\Delta t}{\Delta x(\Delta x + \langle S_1 \rangle - S_0)} ,$$

$$C(M,M+1) = - \frac{2\Delta t}{(\langle S_1 \rangle - S_0)(\Delta x + \langle S_1 \rangle - S_0)} ,$$

$$C(n,n-1) = - \frac{2\Delta t}{(\langle S_{n-M} \rangle - \langle S_{n-M-1} \rangle)(\langle S_{n-M+1} \rangle - \langle S_{n-M-1} \rangle)} ,$$

$$n = M+1, M+2, \dots, M+L(j+1)-1,$$

$$C(n,n) = 1 + \frac{2\Delta t}{(\langle S_{n-M+1} \rangle - \langle S_{n-M} \rangle)(\langle S_{n-M+1} \rangle - \langle S_{n-M-1} \rangle)} \\ + \frac{2\Delta t}{(\langle S_{n-M} \rangle - \langle S_{n-M-1} \rangle)(\langle S_{n-M+1} \rangle - \langle S_{n-M-1} \rangle)} ,$$

$$n = M+1, M+2, \dots, M+L(j+1)-1,$$

$$C(n,n+1) = - \frac{2\Delta t}{(\langle S_{n-M+1} \rangle - \langle S_{n-M} \rangle)(\langle S_{n-M+1} \rangle - \langle S_{n-M-1} \rangle)} ,$$

$$n = M+1, M+2, \dots, M+L(j+1)-2.$$

CHAPTER III

SOME PRELIMINARY RESULTS

The results that are derived in this chapter will be used in Chapter IV to prove convergence.

Lemma 3.1. The sequence of mesh points S_j , $j = 0, 1, 2, \dots, N$, defined by (2.2), is a monotonic nondecreasing sequence of mesh points.

Proof. Since $f, \phi \geq 0$ and $U_{M+L(j),j} = 0$ the maximum principle [13] implies that $U_{i,j} \geq 0$ for $i = 0, 1, 2, \dots, M+L(j)$, $j = 0, 1, 2, \dots, N$. Therefore we have

$$(3.1) \quad \frac{S_1 - S_0}{\Delta t} = \frac{U_{M-1,0}}{\Delta x} = \frac{\phi(b - \Delta x)}{\Delta x} \geq 0$$

which implies that

$$S_1 \geq S_0.$$

Assume that

$$S_j \geq S_{j-1} \geq S_{j-2} \geq \dots \geq S_0.$$

Then

$$(3.2) \quad \frac{S_{j+1} - S_j}{\Delta t} = \frac{U_{M+L(j)-1,j}}{S_j - [S_{j-1}]} \geq 0$$

and from this we conclude that

$$S_{j+1} \geq S_j$$

for $j = 0, 1, 2, \dots, N-1$, which is the desired result.

Lemma 3.2. (a) Let u be a solution of (1.1) where s is a monotonic nondecreasing function. Then there exists a positive

constant A depending only on b ,

$$B = \max \left\{ \max_{0 \leq t \leq T} f(t), \max_{0 \leq x \leq b} \phi(x) \right\}$$

and the Lipschitz constant c in (1.3), such that

$$0 \leq \frac{u(s(t)-\rho, t)}{\rho} \leq A$$

for all $0 \leq t \leq T$ and $0 < \rho < b$.

(b) Let $(S_j, U_{i,j})$ be the solution of (2.1) - (2.3). Then

$$0 \leq \frac{U_{i,j}}{S_j - x_i} \leq A$$

for $0 \leq j \leq N$, $0 \leq i < M+L(j)$, where A is the same as in part (a).

Proof. (a) See Cannon and Hill [1, Lemma 2, page 3].

(b) The maximum principle implies that $U_{i,j} \geq 0$ since $f, \phi \geq 0$ and $U_{M+L(j),j} = 0$. Let

$$A = \max \{Bb^{-1}, c\}.$$

We define

$$W_{i,j}(j_0) \equiv A(S_{j_0} - x_i)$$

for each j_0 , where $0 \leq j_0 \leq N$. It is easy to see that $W_{i,j}(j_0)$ satisfies (2.1) and that

$$W_{0,j}(j_0) \geq f.$$

This follows because $W_{0,j}(j_0) = AS_{j_0} \leq Ab \leq B \leq f$ and $0 < b = S_0$

$\leq S_{j_0}$. For $j = 0$ we have

$$W_{i,0}(j_0) \geq \phi$$

because

$$A(S_{j_0} - x_i) \geq A(b - x_i) \geq C(b - x_i) \geq \phi.$$

For $0 \leq j \leq j_0$ and W on the right-hand boundary,

$$W_{M+L(j),j}^{(j_0)} = A(S_{j_0} - S_j) \geq 0.$$

Since $W_{i,j}^{(j_0)} \geq U_{i,j}$ on the pertinent parts of the boundary, the maximum principle implies that $W_{i,j}^{(j_0)} \geq U_{i,j}$ in the interior, which implies that

$$U_{i,j} \leq W_{i,j}^{(j_0)}$$

for $0 \leq j \leq j_0$, $0 \leq i < M + L(j)$. Thus we can conclude that

$$0 \leq U_{i,j} \leq A(S_j - x_i)$$

for $0 \leq j \leq N$, $0 \leq i < M + L(j)$. This completes the proof of Lemma 3.2. Note that the proof given here parallels that for part (a).

Lemma 3.3. Let $(S_j, U_{i,j})$ be the solution of (2.1) - (2.3). Then there exists a positive constant A such that

$$0 \leq \frac{S_{j+1} - S_j}{\Delta t} \leq A$$

for all $0 \leq j \leq N - 1$.

Proof. Using (3.1), (3.2) and Lemma 3.2, we obtain

$$0 \leq \frac{S_1 - S_0}{\Delta t} = \frac{U_{M-1,0}}{\Delta x} \leq A$$

and

$$0 \leq \frac{S_{j+1} - S_j}{\Delta t} = \frac{U_{M+L(j)-1,j}}{S_j - [S_{j-1}]} \leq A$$

for $j = 1, 2, \dots, N - 1$. This gives us the desired result.

Lemma 3.4. Let

$$x_i = i\Delta x \quad \text{for } i = 0, 1, 2, \dots, M,$$

$$x_{M+i} = \langle S_i \rangle \quad \text{for } i = 0, 1, 2, \dots, L(N),$$

$$\Delta x_i = x_{i+1} - x_i,$$

$$\overline{\Delta x}_i = \begin{cases} \frac{\Delta x_0}{2} & \text{if } i = 0, \\ \frac{1}{2}(\Delta x_i + \Delta x_{i-1}) & \text{if } 0 < i < M, 0 \leq j < J + 1, \\ \frac{\Delta x_{M-1}}{2} & \text{if } i = M, 0 \leq j < J + 1, \\ \frac{1}{2}(\Delta x_i + \Delta x_{i-1}) & \text{if } 0 < i < M + L(j), J + 1 \leq j \leq N, \\ \frac{\Delta x_{M+L(j)-1}}{2} & \text{if } i = M + L(j), J + 1 \leq j \leq N, \end{cases}$$

$$P(M) = \min \{k | S_k \neq S_0\}.$$

$$P(i) = \min \{k | S_k \neq S_{P(i-1)}, k > P(i-1)\} \text{ for } i \geq M + 1,$$

$$H(0) = \max \{k | S_k = S_0\},$$

$$H(j) = \max \{k | S_k = S_{H(j-1)+1}\} \text{ for } j \geq 1.$$

Then it follows that

$$(3.3) \quad \begin{aligned} & \sum_{i=0}^{M-1} \sum_{j=1}^K \left\{ \frac{F_{i+1,j} - F_{i,j}}{\Delta x_i} - \frac{G_{i,j} - G_{i,j-1}}{\Delta t} \right\} \overline{\Delta x}_i \Delta t \\ & + \sum_{i=M}^{M+L(K)-1} \sum_{j=P(i)}^K \left\{ \frac{F_{i+1,j} - F_{i,j}}{\Delta x_i} - \frac{G_{i,j} - G_{i,j-1}}{\Delta t} \right\} \overline{\Delta x}_i \Delta t \end{aligned}$$

$$\begin{aligned}
&= \sum_{j=1}^K \{F_{M+L(j),j} - F_{0,j}\} \Delta t - \sum_{i=0}^{M-1} \{G_{i,K} - G_{i,0}\} \overline{\Delta x}_i \\
&\quad - \sum_{j=0}^{L(K)-1} \{G_{M+j,K} - G_{M+j,H(j)}\} \overline{\Delta x}_{M+j}
\end{aligned}$$

for all $0 < K \leq N$ and arbitrary mesh functions $F_{i,j}$ and $G_{i,j}$.

Proof.

$$\begin{aligned}
&\sum_{i=0}^{M-1} \sum_{j=1}^K \left\{ \frac{F_{i+1,j} - F_{i,j}}{\overline{\Delta x}_i} - \frac{G_{i,j} - G_{i,j-1}}{\Delta t} \right\} \overline{\Delta x}_i \Delta t \\
&\quad + \sum_{i=M}^{M+L(K)-1} \sum_{j=P(i)}^K \left\{ \frac{F_{i+1,j} - F_{i,j}}{\overline{\Delta x}_i} - \frac{G_{i,j} - G_{i,j-1}}{\Delta t} \right\} \overline{\Delta x}_i \Delta t \\
&= \sum_{j=1}^K \sum_{i=0}^{M-1} \{F_{i+1,j} - F_{i,j}\} \Delta t - \sum_{i=0}^{M-1} \sum_{j=1}^K \{G_{i,j} - G_{i,j-1}\} \overline{\Delta x}_i \\
&\quad + \sum_{i=M}^{M+L(K)-1} \sum_{j=P(i)}^K \{F_{i+1,j} - F_{i,j}\} \Delta t \\
&\quad - \sum_{i=M}^{M+L(K)-1} \sum_{j=P(i)}^K \{G_{i,j} - G_{i,j-1}\} \overline{\Delta x}_i \\
&= \sum_{j=1}^K \{F_{M,j} - F_{0,j}\} \Delta t - \sum_{i=0}^{M-1} \{G_{i,K} - G_{i,0}\} \overline{\Delta x}_i \\
&\quad - \sum_{i=M}^{M+L(K)-1} \{G_{i,K} - G_{i,P(i)-1}\} \overline{\Delta x}_i + \sum_{j=P(M)}^K \{F_{M+1,j} - F_{M,j}\} \Delta t \\
&\quad + \sum_{j=P(M+1)}^K \{F_{M+2,j} - F_{M+1,j}\} \Delta t + \sum_{j=P(M+2)}^K \{F_{M+3,j} - F_{M+2,j}\} \Delta t \\
&\quad + \cdots + \{F_{M+L(K),K} - F_{M+L(K)-1,K}\} \Delta t
\end{aligned}$$

$$\begin{aligned}
&= \sum_{j=1}^K \{F_{M,j} - F_{0,j}\} \Delta t - \sum_{i=0}^{M-1} \{G_{i,K} - G_{i,0}\} \overline{\Delta x}_i \\
&\quad - \sum_{j=0}^{L(K)-1} \{G_{M+j,K} - G_{M+j,P(M+j)-1}\} \overline{\Delta x}_{M+j} \\
&\quad + \sum_{j=P(M)}^K \{F_{M+1,j} - F_{M,j}\} \Delta t + \sum_{j=P(M+1)}^K \{F_{M+2,j} - F_{M+1,j}\} \Delta t \\
&\quad + \sum_{j=P(M+2)}^K \{F_{M+3,j} - F_{M+2,j}\} \Delta t + \cdots + \{F_{M+L(K),K} - F_{M+L(K)-1,K}\} \Delta t \\
&= \sum_{j=1}^K \{F_{M+L(j),j} - F_{0,j}\} \Delta t \\
&\quad - \sum_{i=0}^{M-1} \{G_{i,K} - G_{i,0}\} \overline{\Delta x}_i - \sum_{j=0}^{L(K)-1} \{G_{M+j,K} - G_{M+j,H(j)}\} \overline{\Delta x}_{M+j}.
\end{aligned}$$

Thus we have obtained the desired result.

Lemma 3.5. Let $(S_j, U_{i,j})$ be the solution of (2.1) - (2.3).

Then

$$\begin{aligned}
&\sum_{j=1}^K \{S_j \frac{S_{j+1} - S_j}{\Delta t}\} \Delta t - \sum_{j=1}^K f(t_j) \Delta t - \sum_{i=0}^{M-1} \{x_i \phi(x_i)\} \overline{\Delta x}_i \\
&\quad + \sum_{i=0}^{M+L(K)-1} \{x_i U_{i,K}\} \overline{\Delta x}_i = 0
\end{aligned}$$

(3.4)

for all $0 < K \leq N$.

Proof. Assume $0 < K \leq N$ and let

$$\nabla_x U_{i,j} = \frac{U_{i,j} - U_{i-1,j}}{\Delta x_{i-1}},$$

$$\nabla_t U_{i,j} = \frac{U_{i,j} - U_{i,j-1}}{\Delta t},$$

$$\nabla_x U_{i,j} = \frac{U_{i+1,j} - U_{i,j}}{\Delta x_i},$$

$$\delta_x^2 U_{i,j} = \frac{\frac{U_{i+1,j} - U_{i,j}}{\Delta x_i} - \frac{U_{i,j} - U_{i-1,j}}{\Delta x_{i-1}}}{\overline{\Delta x_i}}.$$

we set

$$F_{i,j} = V_{i,j} \nabla_x U_{i,j} - U_{i,j} \nabla_x V_{i,j},$$

$$G_{i,j} = U_{i,j} V_{i,j}$$

in (3.3) to obtain the identity

$$\begin{aligned} & \sum_{i=0}^{M-1} \sum_{j=1}^K \{V_{i,j} \delta_x^2 U_{i,j} - U_{i,j} \delta_x^2 V_{i,j} - V_{i,j} \nabla_t U_{i,j} \\ & \quad - U_{i,j-1} \nabla_t V_{i,j}\} \overline{\Delta x_i} \Delta t \\ & + \sum_{i=M}^{M+L(K)-1} \sum_{j=P(i)}^K \{V_{i,j} \delta_x^2 U_{i,j} - U_{i,j} \delta_x^2 V_{i,j} - V_{i,j} \nabla_t U_{i,j} \\ & \quad - U_{i,j-1} \nabla_t V_{i,j}\} \overline{\Delta x_i} \Delta t \\ (3.5) \quad & + \sum_{j=1}^K \{V_{M+L(j),j} \nabla_x U_{M+L(j),j} - U_{M+L(j),j} \nabla_x V_{M+L(j),j} - V_{0,j} \nabla_x U_{0,j} \end{aligned}$$

$$\begin{aligned}
& + U_{0,j} \nabla_x V_{0,j} \Delta t - \sum_{i=0}^{M-1} \{U_{i,K} V_{i,K} - U_{i,0} V_{i,0}\} \overline{\Delta x}_i \\
& - \sum_{j=0}^{L(K)-1} \{U_{M+j,K} V_{M+j,K} - U_{M+j,H(j)} V_{M+j,H(j)}\} \overline{\Delta x}_i.
\end{aligned}$$

Next we let $U_{i,j}$ be the unique solution of (2.1) - (2.3) and we set $V_{i,j} = -x_i$ in (3.5). Then the left hand side of (3.5) is zero. Since $U_{M+L(j),j} = 0$, $U_{M+j,H(j)} = 0$, $V_{0,j} = 0$, (3.5) becomes

$$\begin{aligned}
& \sum_{j=1}^K \{S_j \frac{S_{j+1} - S_j}{\Delta t}\} \Delta t - \sum_{j=1}^K f(t_j) \Delta t - \sum_{i=0}^{M-1} \{x_i \phi(x_i)\} \overline{\Delta x}_i \\
& + \sum_{i=0}^{M+L(K)-1} \{x_i U_{i,K}\} \overline{\Delta x}_i = 0,
\end{aligned}$$

which completes the proof of Lemma 3.5.

Lemma 3.6. Let $(S_j, U_{i,j})$ be the solution of (2.1) - (2.3).

Then

$$\begin{aligned}
& \frac{1}{2}(S_{K+1}^2 - S_1^2) + O(\Delta t) - \sum_{j=1}^K f(t_j) \Delta t - \sum_{i=1}^{M-1} \{x_i \phi(x_i)\} \overline{\Delta x}_i \\
(3.6) \quad & - x_0 \phi(x_0) \frac{\Delta x_0}{2} - x_M \phi(x_M) \frac{\Delta x_{M-1}}{2} + \sum_{i=1}^{M+L(K)-1} \{x_i U_{i,K}\} \overline{\Delta x}_i \\
& + x_0 U_{0,K} \frac{\Delta x_0}{2} + x_{M+L(K)} U_{M+L(K),K} \frac{\Delta x_{M+L(K)-1}}{2} = 0
\end{aligned}$$

for all $0 < K \leq N$. We note that S_{N+1} is defined since our finite difference solution $(S_j, U_{i,j})$ is defined for $j = 0, 1, 2, \dots, N$, $i = 0, 1, 2, \dots, M+L(N)$.

Proof. Note that for $0 < K \leq N$ we can write the sum

$$\sum_{j=1}^K \left\{ S_j \frac{S_{j+1} - S_j}{\Delta t} \right\} \Delta t$$

in the following way

$$\begin{aligned} \sum_{j=1}^K \left\{ S_j \frac{S_{j+1} - S_j}{\Delta t} \right\} \Delta t &= \sum_{j=1}^K \left\{ S_j \frac{S_{j+1} - S_j}{\Delta t} + \frac{(S_{j+1} + S_j)(S_{j+1} - S_j)}{2\Delta t} \right. \\ &\quad \left. - \frac{(S_{j+1} + S_j)(S_{j+1} - S_j)}{2\Delta t} \right\} \Delta t \\ (3.7) \quad &= \sum_{j=1}^K \left\{ \left[S_j - \frac{(S_{j+1} + S_j)}{2} \right] \frac{S_{j+1} - S_j}{\Delta t} + \frac{(S_{j+1} + S_j)(S_{j+1} - S_j)}{2\Delta t} \right\} \Delta t \\ &= \sum_{j=1}^K \left\{ \frac{S_{j+1}^2 - S_j^2}{2\Delta t} - \frac{(S_{j+1} - S_j)^2}{2\Delta t} \right\} \Delta t \\ &= \frac{1}{2}(S_{K+1}^2 - S_1^2) - \frac{1}{2} \sum_{j=1}^K \left\{ \frac{(S_{j+1} - S_j)^2}{(\Delta t)^2} \right\} (\Delta t)^2. \end{aligned}$$

Using Lemma 3.3 we obtain

$$\begin{aligned} \left| \frac{1}{2} \sum_{j=1}^K \left\{ \frac{(S_{j+1} - S_j)^2}{(\Delta t)^2} \right\} (\Delta t)^2 \right| &\leq \frac{1}{2} K A^2 (\Delta t)^2 \\ &= \frac{1}{2} (K \Delta t) (A^2 \Delta t) \\ &\leq \frac{1}{2} T A^2 \Delta t, \end{aligned}$$

which implies

$$(3.8) \quad \frac{1}{2} \sum_{j=1}^K \left\{ \frac{(S_{j+1} - S_j)^2}{(\Delta t)^2} \right\} (\Delta t)^2 = O(\Delta t).$$

Combining (3.7) and (3.8) we get

$$(3.9) \quad \sum_{j=1}^K \left\{ S_j \frac{S_{j+1} - S_j}{\Delta t} \right\} \Delta t = \frac{1}{2} (S_{K+1}^2 - S_1^2) + O(\Delta t).$$

Next, recalling that $\phi(x_M) = 0$, we note that

$$(3.10) \quad \sum_{i=0}^{M-1} \{x_i \phi(x_i)\} \overline{\Delta x}_i = \sum_{i=1}^{M-1} \{x_i \phi(x_i)\} \overline{\Delta x}_i + x_0 \phi(x_0) \frac{\Delta x_0}{2} \\ + x_M \phi(x_M) \frac{\Delta x_{M-1}}{2}.$$

Furthermore,

$$(3.11) \quad \sum_{i=0}^{M+L(K)-1} \{x_i U_{i,K}\} \overline{\Delta x}_i = \sum_{i=1}^{M+L(K)-1} \{x_i U_{i,K}\} \overline{\Delta x}_i + x_0 U_{0,K} \frac{\Delta x_0}{2} \\ + x_{M+L(K)} U_{M+L(K),K} \frac{\Delta x_{M+L(K)-1}}{2},$$

using that $U_{M+L(K),K} = 0$.

Combining (3.4), (3.9), (3.10) and (3.11), we obtain the identity

$$\frac{1}{2} (S_{K+1}^2 - S_1^2) + O(\Delta t) - \sum_{j=1}^K f(t_j) \Delta t - \sum_{i=1}^{M-1} \{x_i \phi(x_i)\} \overline{\Delta x}_i - x_0 \phi(x_0) \frac{\Delta x_0}{2} \\ - x_M \phi(x_M) \frac{\Delta x_{M-1}}{2} + \sum_{i=1}^{M+L(K)-1} \{x_i U_{i,K}\} \overline{\Delta x}_i + x_0 U_{0,K} \frac{\Delta x_0}{2}$$

$$+ x_{M+L(K)}^U \frac{\Delta x_{M+L(K)-1}}{2} = 0,$$

which is (3.6), and we are done.

CHAPTER IV

CONVERGENCE

Let $\{(\Delta x^n, \Delta t^n)\}_{n=1}^{\infty}$ be a sequence of ordered pairs, where $\{\Delta x^n\}_{n=1}^{\infty}$ and $\{\Delta t^n\}_{n=1}^{\infty}$ are nonincreasing sequences of real numbers both of which approach zero as $n \rightarrow \infty$. For each member of the sequence $\{(\Delta x^n, \Delta t^n)\}_{n=1}^{\infty}$ we obtain the corresponding S_j^n 's from the scheme described by (2.1) - (2.3). Next we define a continuous function $s^n(t)$ using piecewise linear interpolation. Thus we have a sequence of functions $\{s^n\}_{n=1}^{\infty}$ defined for $0 \leq t \leq T$ and such that $s^n(t_j^n) = S_j^n$.

The following notation will be used later. It corresponds to earlier notation, only now involves the sequence $\{(\Delta x^n, \Delta t^n)\}_{n=1}^{\infty}$

$$x_i^n = i\Delta x^n \quad \text{for } i = 0, 1, 2, \dots, M,$$

$$x_{M+i}^n = \langle S_i^n \rangle \quad \text{for } i = 0, 1, 2, \dots, L(N),$$

$$\Delta x_i^n = x_{i+1}^n - x_i^n,$$

$$\overline{\Delta x}_i^n = \begin{cases} \frac{\Delta x_0^n}{2} & \text{if } i = 0, \\ \frac{1}{2}(\Delta x_i^n + \Delta x_{i-1}^n) & \text{if } 0 < i < M, \quad 0 \leq j < J + 1, \\ \frac{\Delta x_{M-1}^n}{2} & \text{if } i = M, \quad 0 \leq j < J + 1, \\ \frac{1}{2}(\Delta x_i^n + \Delta x_{i-1}^n) & \text{if } 0 < i < M + L(j), \quad J + 1 \leq j \leq N, \\ \frac{\Delta x_{M+L(j)-1}^n}{2} & \text{if } i = M + L(j), \quad J + 1 \leq j \leq N, \end{cases}$$

$$t_j^n = j\Delta t^n \quad \text{for } j = 0, 1, \dots, N.$$

Also, let $(S_j^n, U_{i,j}^n)$ be the solution of (2.1) - (2.3) for the mesh spacing $(\Delta x^n, \Delta t^n)$, let u be the solution of (1.1), and define $u_{i,j}^n = u(x_i^n, t_j^n)$.

Next we prove the following lemma concerning the sequence $\{s^n\}_{n=1}^\infty$:

Lemma 4.1. The sequence $\{s^n\}_{n=1}^\infty$ is equicontinuous and uniformly bounded on $[0, T]$.

Proof. Clearly each s^n is uniformly continuous on the interval $[0, T]$. To show that the sequence $\{s^n\}_{n=1}^\infty$ is equicontinuous it suffices to show that each s^n is Lipschitz continuous with the same Lipschitz constant A .

First we note that using Lemma 3.3 and the fact that $s^n(t_j^n) = S_j^n$ we have

$$(4.1) \quad |s^n(t_j^n) - s^n(t_{j-1}^n)| \leq A \Delta t^n$$

for $j = 1, 2, \dots, N$. Using (4.1) and the linearity of s^n on each interval, it is clear that

$$(4.2) \quad |s^n(\bar{t}) - s^n(\bar{t})| \leq A |\bar{t} - \bar{t}|$$

for $\bar{t}, \bar{t} \in [t_{j-1}^n, t_j^n]$ and $j = 1, 2, \dots, N$.

Next we let $t', t'' \in [0, T]$, but possibly fall in different subintervals of $[0, T]$. Assume t' is in the subinterval $[t_{\ell-1}^n, t_\ell^n]$ and t'' in $[t_{k-1}^n, t_k^n]$, where ℓ and k may be different. Without loss of generality we may assume that $k \geq \ell$. Using (4.1) and (4.2) we can

conclude that

$$\begin{aligned}
 |s^n(t'') - s^n(t')| &\leq |s^n(t'') - s^n(t_{k-1}^n)| + |s^n(t_{k-1}^n) - s^n(t_{k-2}^n)| \\
 &\quad + \cdots + |s^n(t_{\ell+1}^n) - s^n(t_{\ell}^n)| + |s^n(t_{\ell}^n) - s^n(t')| \\
 &\leq A|t'' - t_{k-1}^n| + A|t_{k-1}^n - t_{k-2}^n| + \cdots \\
 &\quad + A|t_{\ell+1}^n - t_{\ell}^n| + A|t_{\ell}^n - t'| \\
 &= A|t'' - t'|.
 \end{aligned}$$

Thus each s^n is Lipschitz continuous with Lipschitz constant A .

To prove uniform boundedness we use Lemma 3.3 to obtain

$$\begin{aligned}
 S_N^n &\leq S_{N-1}^n + A\Delta t^n \\
 &\leq S_{N-2}^n + 2A\Delta t^n \\
 &\leq \cdots \leq S_0^n + N\Delta t^n A \\
 &\leq b + TA,
 \end{aligned}$$

which together with Lemma 3.1, the fact that $S_0^n = b > 0$, and the definition of s^n implies that

$$0 < s^n(t) \leq b + TA$$

for $t \in [0, T]$ and all n . Therefore the sequence of functions $\{s^n\}_{n=1}^{\infty}$

is uniformly bounded. Thus the proof of Lemma 4.1 is complete.

By Lemma 4.1, the sequence of functions $\{s^n\}_{n=1}^{\infty}$ satisfies the hypotheses of Arzela's Theorem. Hence there exists a uniformly convergent subsequence of the sequence $\{s^n\}_{n=1}^{\infty}$. We denote this subsequence by $\{s^n\}_{n=1}^{\infty}$ and its limit by s . It will be shown later that the selection of subsequences is unnecessary.

Since the subsequence $\{s^n\}_{n=1}^{\infty}$ converges uniformly to s and each s^n is continuous, clearly, then, it follows that s is continuous.

Furthermore we can establish the following lemmas concerning s :

Lemma 4.2. s is monotone nondecreasing.

Proof. Since s^n converges uniformly to s and each s^n is monotone nondecreasing we can write

$$\begin{aligned} s(t_1) - s(t_2) &= [s(t_1) - s^n(t_1)] + [s^n(t_1) - s^n(t_2)] \\ (4.3) \qquad &+ [s^n(t_2) - s(t_2)] \leq \frac{\varepsilon}{2} + 0 + \frac{\varepsilon}{2} = \varepsilon \end{aligned}$$

for n sufficiently large and $t_1 \leq t_2$, where ε is an arbitrary positive number. Therefore (4.3) implies that $s(t_1) - s(t_2) \leq 0$ since ε is arbitrary. Hence s is monotone nondecreasing.

Lemma 4.3. s is Lipschitz continuous.

Proof. In the proof of Lemma 4.1 we saw that each s^n is Lipschitz continuous with Lipschitz constant A . Let $t', t'' \in [0, T]$. We note that

$$\begin{aligned}
(4.4) \quad & |s(t') - s(t'')| \leq |s(t') - s^n(t')| \\
& + |s^n(t') - s^n(t'')| + |s^n(t'') - s(t'')|.
\end{aligned}$$

Since $\{s^n\}_{n=1}^\infty$ converges uniformly to s and each s^n is Lipschitz continuous with Lipschitz constant A , for n sufficiently large, (4.4) becomes

$$(4.5) \quad |s(t') - s(t'')| \leq \varepsilon + A|t' - t''|,$$

where ε is an arbitrary positive number that approaches zero as n approaches infinity. Finally letting n go to infinity in (4.5), we see that

$$|s(t') - s(t'')| \leq A|t' - t''|$$

for $t', t'' \in [0, T]$, which is the desired result.

Now we let u denote the unique solution of (1.1), where s is the uniform limit of the subsequence $\{s^n\}_{n=1}^\infty$. Since f and ϕ are continuously differentiable and s is Lipschitz continuous, we know that a unique solution of (1.1) exists by Hill and Kotlow [8, Lemma 2.1, page 65].

We take $u_{i,j}^n$ and $U_{i,j}^n$ to be as defined previously. Further, we define them to be zero outside their respective domains. Also we set $s_\varepsilon(t) = s(t) - \varepsilon$ for $0 \leq t \leq T$, where ε is a small arbitrary positive number. Furthermore let us define $D_\varepsilon = \{(x, t) \mid 0 < x < s_\varepsilon(t), 0 < t \leq T\}$ and $D_n = \{(x, t) \mid 0 < x < s^n(t), 0 < t \leq T\}$. $\overline{D}_\varepsilon$ and $\overline{D}_n$ will

represent the closures of D_ε and D_n , respectively. Let Ω be an arbitrary domain. Then define $C^{2,1}(\Omega) = \{u | u \in C^2 \text{ with respect to } x \text{ and } u \in C^1 \text{ with respect to } t \text{ for } (x,t) \in \Omega\}$.

The next lemma will be needed in the proof of Theorem 4.1.

Lemma 4.4. Under the assumptions on f and ϕ in Chapter I, let u be the unique solution of (1.1) with s the uniform limit of the subsequence $\{s^n\}_{n=1}^\infty$. Then $u \in C^{2,1}(\bar{D}_\varepsilon)$.

Proof. From Hill and Kotlow [8, Lemma 2.1, page 65] we have that $u \in C^{2,1}(D)$ for $D = \{(x,t) | 0 < x < s(t), 0 < t \leq T\}$. Application of the techniques of Gevrey [7] to $R = \{(x,t) | 0 \leq x \leq b, 0 \leq t \leq T\}$ further shows that $u \in C^{2,1}(R)$. Combining these two results, then, we have that $u \in C^{2,1}(\bar{D}_\varepsilon)$ and we are done.

Note that there is a subsequence of the entire sequence $\{(\Delta x^n, \Delta t^n)\}_{n=1}^\infty$ corresponding to the subsequence $\{s^n\}_{n=1}^\infty$. We will also denote it by $\{(\Delta x^n, \Delta t^n)\}_{n=1}^\infty$. Clearly, just as for the entire sequence, for the subsequence, as $n \rightarrow \infty$, Δx^n and $\Delta t^n \rightarrow 0$. Let $U_{i,j}^n$ be as defined previously and let $\tilde{U}_{i,j}^n$ be the same only based on the subsequence $\{(\Delta x^n, \Delta t^n)\}_{n=1}^\infty$. Let $u_{i,j}^n$ be as before. Whether $u_{i,j}^n$ corresponds to the complete sequence or the subsequence will be clear from context.

We say that $U_{i,j}^n$ ($\tilde{U}_{i,j}^n$) converges uniformly with respect to i,j to u on the region $D_\varepsilon = \{(x,t) | 0 \leq x < \infty, 0 \leq t \leq T\}$ as $n \rightarrow \infty$, if given $\varepsilon > 0$, then $\max_{i,j} |U_{i,j}^n - u_{i,j}^n| < \varepsilon$ ($\max_{i,j} |\tilde{U}_{i,j}^n - u_{i,j}^n| < \varepsilon$) for n sufficiently large. We note that as $n \rightarrow \infty$, $\Delta x_i^n \rightarrow 0$ for $i = 0, 1, 2, \dots, M-1$ by the definition of the sequence $\{(\Delta x^n, \Delta t^n)\}_{n=1}^\infty$ and $\Delta x_i^n \rightarrow 0$ for $i = M, M+1, M+2, \dots$ as a result of Lemma 3.3.

We are now ready to prove the following convergence theorem:

Theorem 4.1. Let $(S_j^n, \tilde{U}_{i,j}^n)$ be the solution of (2.1) - (2.3), and let u be the solution of (1.1) with s the uniform limit of the subsequence $\{s^n\}_{n=1}^\infty$. Then $\tilde{U}_{i,j}^n$ converges uniformly with respect to i, j to u on D_∞ as $n \rightarrow \infty$.

Proof. Let us take $z_{i,j}^n = u_{i,j}^n - \tilde{U}_{i,j}^n$. Since $u_{i,j}^n$ satisfies (1.1) and $\tilde{U}_{i,j}^n$ satisfies (2.1) - (2.3), it follows, using the standard approach, that $z_{i,j}^n$ satisfies

$$(4.6) \quad \frac{z_{i,j}^n - z_{i,j-1}^n}{\Delta t^n} = \frac{\frac{z_{i+1,j}^n - z_{i,j}^n}{\Delta x_i^n} - \frac{z_{i,j}^n - z_{i-1,j}^n}{\Delta x_{i-1}^n}}{\frac{1}{2}(\Delta x_i^n + \Delta x_{i-1}^n)} + Q_n(x_i^n, t_j^n)$$

in $D_\varepsilon \cap D_n$ and $Q_n(x_i^n, t_j^n)$ is given by

$$Q_n(x_i^n, t_j^n) = Q'_n(x_i^n, t_j^n) + Q''_n(x_i^n, t_j^n),$$

where

$$Q'_n(x_i^n, t_j^n) = \frac{u_{i,j}^n - u_{i,j-1}^n}{\Delta t^n} - \frac{\partial u(x_i^n, t_j^n)}{\partial t},$$

$$Q''_n(x_i^n, t_j^n) = \frac{\partial^2 u(x_i^n, t_j^n)}{\partial x^2} - \frac{\frac{u_{i+1,j}^n - u_{i,j}^n}{\Delta x_i^n} - \frac{u_{i,j}^n - u_{i-1,j}^n}{\Delta x_{i-1}^n}}{\frac{1}{2}(\Delta x_i^n + \Delta x_{i-1}^n)}.$$

Seeing that this definition applies on D_ε as well as $D_\varepsilon \cap D_n$ we

consider Q_n defined on D_ε . Using that $u \in C^{2,1}(\overline{D_\varepsilon})$, (see Lemma 4.4), we can easily show that Q_n approaches zero uniformly in D_ε with respect to x_i^n and t_j^n as $n \rightarrow \infty$. Here uniform convergence is understood in a sense analogous to that for $U_{i,j}^n$ and $\tilde{U}_{i,j}^n$. Note that (4.6) can be written in the following form:

$$(4.7) \quad \begin{aligned} z_{i,j}^n = & z_{i,j-1}^n + \frac{2\Delta t^n}{(\Delta x_i^n + \Delta x_{i-1}^n)} \left[\frac{1}{\Delta x_i^n} (z_{i+1,j}^n - z_{i,j}^n) \right. \\ & \left. - \frac{1}{\Delta x_{i-1}^n} (z_{i,j}^n - z_{i-1,j}^n) \right] + \Delta t^n Q_n(x_i^n, t_j^n). \end{aligned}$$

For points (x_i^n, t_j^n) such that $\min\{s_\varepsilon(t_j^n), s^n(t_j^n)\} \leq x_i^n \leq \max\{s^n(t_j^n), s(t_j^n)\}$, we obtain, using Lemma 3.2, the following

inequalities:

$$0 \leq u_{i,j}^n \leq A\varepsilon + A|s^n(t_j^n) - s(t_j^n)|,$$

$$0 \leq \tilde{U}_{i,j}^n \leq A\varepsilon + A|s^n(t_j^n) - s(t_j^n)|.$$

These imply that

$$|z_{i,j}^n| \leq A\varepsilon + A|s^n(t_j^n) - s(t_j^n)|$$

for points in the region indicated.

We now consider the whole region D_∞ by defining mesh points outside D_n as follows: $x_{M+L(N)+i}^n = x_{M+L(N)}^n + i\Delta x^n$ for $i = 0, 1, 2, \dots$, $t_j^n = t_j^n$, $j = 0, 1, 2, \dots, N$. For each j , $j = 0, 1, 2, \dots, N$, we note that either $\max_{i \geq 0} z_{i,j}^n = 0$, $\max_{i \geq 0} z_{i,j}^n \leq A\epsilon + A|s^n(t_j^n) - s(t_j^n)|$, or $\max_{i \geq 0} z_{i,j}^n$ occurs in $D_\epsilon \cap D_n$.

Assume that the $\max_{i \geq 0} z_{i,j}^n$ occurs at a point (x_i^n, t_j^n) in $D_\epsilon \cap D_n$.

Using (4.7) we obtain

$$\begin{aligned} z_{i,j}^n = z_{i,j-1}^n &+ \frac{2\Delta t^n}{(\Delta x_i^n + \Delta x_{i-1}^n)} \left[\frac{1}{\Delta x_i^n} (z_{i+1,j}^n - z_{i,j}^n) \right. \\ &\left. - \frac{1}{\Delta x_{i-1}^n} (z_{i,j}^n - z_{i-1,j}^n) \right] + \Delta t^n Q_n(x_i^n, t_j^n). \end{aligned}$$

Since

$$z_{i+1,j}^n - z_{i,j}^n \leq 0,$$

$$z_{i,j}^n - z_{i-1,j}^n \geq 0,$$

it follows that

$$(4.8) \quad z_{i,j}^n \leq z_{i,j-1}^n + \Delta t^n Q_n(x_i^n, t_j^n)$$

$$\leq \max_{i \geq 0} z_{i,j-1}^n + \Delta t \max_{\substack{(x_k^n, t_p^n) \in D_\varepsilon \\ (x_k^n, t_p^n) \in D_n}} Q_n(x_k^n, t_p^n).$$

Similarly it follows that

$$(4.9) \quad z_{i,j}^n \geq \min_{i \geq 0} z_{i,j-1}^n + \Delta t \min_{\substack{(x_k^n, t_p^n) \in D_\varepsilon \\ (x_k^n, t_p^n) \in D_n}} Q_n(x_k^n, t_p^n)$$

if we assume that the $\min_{i \geq 0} z_{i,j}^n$ occurs at a point (x_i^n, t_j^n) in $D_\varepsilon \cap D_n$.

We will use (4.8) and (4.9) in a moment but first, recalling the three possibilities for $\max_{i \geq 0} z_{i,j}^n$ and noting that there are similar possibilities for $\min_{i \geq 0} z_{i,j}^n$, we set

$$M_\mu^n(j) = \max[0, A \varepsilon + A \max_{0 \leq k \leq N} |s_k^n(t_k^n) - s(t_k^n)|, z_{i,j}^n],$$

$$M_\ell^n(j) = \min[0, -A \varepsilon - A \max_{0 \leq k \leq N} |s_k^n(t_k^n) - s(t_k^n)|, z_{i,j}^n].$$

Then it follows that

$$M_\ell^n(j) \leq z_{i,j}^n \leq M_\mu^n(j)$$

for $i = 0, 1, 2, \dots; j = 0, 1, 2, \dots, N$. This implies that

$$\max_{i \geq 0} |z_{i,j}^n| \leq \max\{|M_\ell^n(j)|, |M_\mu^n(j)|\}$$

for $j = 0, 1, 2, \dots, N$. Now using (4.8) and (4.9) here, we get

$$\max_{i \geq 0} |z_{i,j}^n| \leq \max\{A \varepsilon + A \max_{0 \leq k \leq N} |s^n(t_k^n) - s(t_k^n)|,$$

(4.10)

$$+ \Delta t^n \max_{(x_k^n, t_p^n) \in D_\varepsilon} |Q_n(x_k^n, t_p^n)|\}.$$

Using induction, inequality (4.10) gives us that

$$\begin{aligned} \max_{i \geq 0} |z_{i,j}^n| &\leq A \varepsilon + A \max_{0 \leq k \leq N} |s^n(t_k^n) - s(t_k^n)| \\ &\quad + j \Delta t^n \max_{(x_k^n, t_p^n) \in D_\varepsilon} |Q_n(x_k^n, t_p^n)| \end{aligned}$$

for $j = 0, 1, 2, \dots, N$, where we have also used the fact that

$$\max_{i \geq 0} |z_{i,0}^n| = 0. \quad \text{Thus}$$

$$\max_{\substack{i \geq 0 \\ 0 \leq j \leq N}} |z_{i,j}^n| \leq A \varepsilon + A \max_{0 \leq k \leq N} |s^n(t_k^n) - s(t_k^n)|$$

(4.11)

$$+ T \max_{(x_k^n, t_p^n) \in D_\varepsilon} |Q_n(x_k^n, t_p^n)|.$$

Let $\varepsilon > 0$ be given. From (4.11), replacing ε there by $\frac{\varepsilon}{3A}$, while recalling that s^n converges uniformly to s and Q_n to zero, we obtain

$$\max_{\substack{i \geq 0 \\ 0 \leq j \leq N}} |z_{i,j}^n| \leq A\left(\frac{\varepsilon}{3A}\right) + A\left(\frac{\varepsilon}{3A}\right) + T\left(\frac{\varepsilon}{3T}\right) = \varepsilon$$

for n sufficiently large. Thus $\tilde{U}_{i,j}^n$ converges uniformly to u on the region D_∞ as $n \rightarrow \infty$; hence the proof of Theorem 4.1 is complete.

Theorem 4.2. Let u be the unique solution of (1.1), where s is the uniform limit of the subsequence $\{s^n\}_{n=1}^\infty$. Then the pair (s, u) is the unique solution of (1.1), (1.2).

Proof. We shall show that (s, u) satisfies (1.4). First we note that

$$(4.12) \quad (S_1^n)^2 = b^2 + O(\Delta t^n)$$

since

$$\begin{aligned} |(S_1^n)^2 - b^2| &= |(S_1^n)^2 - (S_0^n)^2| = |S_1^n - S_0^n| |S_1^n + S_0^n| \\ &\leq A\Delta t^n (2b + A\Delta t^n). \end{aligned}$$

We also note that

$$(4.13) \quad (S_{K+1}^n)^2 = (S_K^n)^2 + O(\Delta t^n)$$

for $K = 1, 2, \dots, N$, since

$$\begin{aligned}
|(S_{K+1}^n)^2 - (S_K^n)^2| &= |S_{K+1}^n - S_K^n| |S_{K+1}^n + S_K^n| \\
&\leq A\Delta t^n (2S_K^n + A\Delta t^n) \\
&\leq A\Delta t^n (2b + 2TA + A\Delta t^n).
\end{aligned}$$

Therefore, using (4.12) and (4.13), (3.6) can be written as

$$\begin{aligned}
&\frac{1}{2}[(S_K^n)^2 - b^2] + o(\Delta t^n) - \sum_{j=1}^K f(t_j^n) \Delta t^n \\
&- \sum_{i=1}^{M-1} \{x_i^n \phi(x_i^n)\} \overline{\Delta x_i^n} - x_0^n \phi(x_0^n) \frac{\Delta x_0^n}{2} \\
(4.14) \quad &- x_M^n \phi(x_M^n) \frac{\Delta x_{M-1}^n}{2} + \sum_{i=1}^{M+L(K)-1} \{x_{i,K}^{\tilde{n}} \tilde{U}_{i,K}^{\tilde{n}}\} \overline{\Delta x_i^n} \\
&+ x_{0,K}^{\tilde{n}} \tilde{U}_{0,K}^{\tilde{n}} \frac{\Delta x_0^n}{2} + x_{M+L(K),K}^{\tilde{n}} \tilde{U}_{M+L(K),K}^{\tilde{n}} \frac{\Delta x_{M+L(K)-1}^n}{2} = 0
\end{aligned}$$

for $0 < K \leq N$.

Next we define

$$I[\tilde{U}^n(K)] = x_{0,K}^{\tilde{n}} \tilde{U}_{0,K}^{\tilde{n}} \frac{\Delta x_0^n}{2} + \sum_{i=1}^{M+L(K)-1} \{x_{i,K}^{\tilde{n}} \tilde{U}_{i,K}^{\tilde{n}}\} \overline{\Delta x_i^n}$$

$$+ x_{M+L(K)}^n \tilde{u}_{M+L(K),K}^n \frac{\Delta x_{M+L(K)}^n - 1}{2} = \sum_{i=0}^{M+L(K)} \{x_{i,K}^n \tilde{u}_{i,K}^n\} \overline{\Delta x_i^n},$$

recalling the definition of $\overline{\Delta x_i^n}$ at the beginning of Chapter IV, and let $I[u^n(K)]$ be defined similarly. Let $0 < t' \leq T$. Then t' must lie in some interval $[(K-1)\Delta t^n, K\Delta t^n]$ for each n . Also we set $u_{i,K}^n = u(x_i^n, t')$, and in the definition of $I[u^n(K')]$, we take $L(K') = L(K)$.

We will now show that $|I[u^n(K)] - I[u^n(K')]| \rightarrow 0$ as $n \rightarrow \infty$. (Note that in this expression and throughout the rest of this proof $K = K(n)$.) First we define $N_1(K) = \{i | 0 \leq x_i < s(t'), 0 \leq i \leq M + L(K)\}$, $N_2(K) = \{i | s(t') \leq x_i < s(t_K^n), M \leq i \leq M + L(K)\}$, and $N_3(K) = \{i | s(t_K^n) \leq x_i, M \leq i \leq M + L(K)\}$. Possibly $N_2(K)$ and $N_3(K)$ may be empty. Assume that $\varepsilon > 0$ is given. Since u is uniformly continuous on $0 \leq x \leq s(t)$, $0 \leq t \leq T$ and s is uniformly continuous on $0 \leq t \leq T$, we can make

$$(4.15) \quad |u_{i,K}^n - u_{i,K'}^n| < \frac{\varepsilon}{2(b+TA)^2}$$

for $i \in N_1(K)$ and

$$(4.16) \quad |s(t_K^n) - s(t')| < \frac{\varepsilon}{2A(b+TA)^2}$$

by taking n sufficiently large. Also, using Lemma 3.2(a) we have

$$(4.17) \quad |u_{i,K}^n| \leq A |s(t_K^n) - x_i^n|.$$

Further, as seen in the proof of Lemma 4.1 we had that $0 \leq S_j^n \leq b + TA$,

thus it follows easily that $0 \leq x_i^n \leq b + TA$ for $0 \leq i \leq M + L(K)$.

Hence for n sufficiently large we obtain

$$\begin{aligned}
 |I[u^n(K)] - I[u^n(K')]| &= \left| \sum_{i \in N_1(K)} x_i^n (u_{i,K}^n - u_{i,K'}^n) \overline{\Delta x_i^n} \right. \\
 &\quad + \sum_{i \in N_2(K)} x_i^n u_{i,K}^n \overline{\Delta x_i^n} \left| \leq \sum_{i \in N_1(K)} x_i^n |u_{i,K}^n - u_{i,K'}^n| \overline{\Delta x_i^n} \right. \\
 &\quad + \sum_{i \in N_2(K)} x_i^n |u_{i,K}^n| \overline{\Delta x_i^n} \leq \sum_{i \in N_1(K)} x_i^n |u_{i,K}^n - u_{i,K'}^n| \overline{\Delta x_i^n} \\
 (4.18) \quad &\quad + \sum_{i \in N_2(K)} x_i^n A |s(t_K^n) - s(t')| \overline{\Delta x_i^n} \leq \sum_{i \in N_1(K)} x_i^n |u_{i,K}^n - u_{i,K'}^n| \overline{\Delta x_i^n} \\
 &\quad + \sum_{i \in N_2(K)} x_i^n A |s(t_K^n) - s(t')| \overline{\Delta x_i^n} \leq \sum_{i \in N_1(K)} (b + TA) \left[\frac{\varepsilon}{2(b+TA)^2} \right] \overline{\Delta x_i^n} \\
 &\quad + \sum_{i \in N_2(K)} A(b + TA) \left[\frac{\varepsilon}{2A(b+TA)^2} \right] \overline{\Delta x_i^n} = \frac{\varepsilon}{2(b+TA)} \sum_{i \in N_1(K)} \overline{\Delta x_i^n} \\
 &\quad + \frac{\varepsilon}{2(b+TA)} \sum_{i \in N_2(K)} \overline{\Delta x_i^n} \leq \frac{\varepsilon}{2(b+TA)} (b + TA) + \frac{\varepsilon}{2(b+TA)} (b + TA) \\
 &\quad = \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon
 \end{aligned}$$

where in the process of the above derivation we have used (4.15), (4.16), and (4.17). We have also used, in going from inequality three to four that $|s(t_K^n) - x_i^n| \leq |s(t_K^n) - s(t')|$ for $i \in N_2(K)$. Furthermore, in going from inequality six to seven we have used the facts

$$\text{that } \sum_{i \in N_1(K)} \overline{\Delta x_i^n} \leq x_{M+L(K)}^n \leq b + TA \text{ and } \sum_{i \in N_2(K)} \overline{\Delta x_i^n} \leq x_{M+L(K)}^n \leq b + TA.$$

Therefore $|I[u^n(K)] - I[u^n(K')]| \rightarrow 0$ as $n \rightarrow \infty$.

Next we prove that $|I[u^n(K')] - \int_0^{s(t')} \xi u(\xi, t') d\xi| \rightarrow 0$ as $n \rightarrow \infty$.

First we define $i^*(K) = \max\{i | x_i \leq s(t'), M \leq i \leq M + L(K)\}$. We note that if $i^*(K) = M + L(K)$ then

$$\begin{aligned} I[u^n(K')] &= x_0^n u_{0,K'}^n \frac{\Delta x_0^n}{2} + \sum_{i=1}^{i^*(K)-1} x_i^n u_{i,K'}^n \overline{\Delta x_i^n} \\ &\quad + x_{i^*(K)}^n u_{i^*(K),K'}^n \frac{\Delta x_{i^*(K)}^n - 1}{2} \leq x_0^n u_{0,K'}^n \frac{\Delta x_0^n}{2} \\ &\quad + \sum_{i=1}^{i^*(K)-1} x_i^n u_{i,K'}^n \overline{\Delta x_i^n} + x_{i^*(K)}^n u_{i^*(K),K'}^n \left[\frac{\Delta x_{i^*(K)}^n - 1}{2} + \frac{\Delta x_{i^*(K)}^n}{2} \right] \end{aligned}$$

However, if $i^*(K) \neq M + L(K)$ we have

$$\begin{aligned} I[u^n(K')] &= x_0^n u_{0,K'}^n \frac{\Delta x_0^n}{2} + \sum_{i=1}^{i^*(K)-1} x_i^n u_{i,K'}^n \overline{\Delta x_i^n} \\ &\quad + x_{i^*(K)}^n u_{i^*(K),K'}^n \left[\frac{\Delta x_{i^*(K)}^n - 1}{2} + \frac{\Delta x_{i^*(K)}^n}{2} \right]. \end{aligned}$$

Let $\varepsilon > 0$ be given. Then for n sufficiently large

$$|I[u^n(K')] - \int_0^{s(t')} \xi u(\xi, t') d\xi| \leq x_0^n u_{0,K'}^n \frac{\Delta x_0^n}{2}$$

$$\begin{aligned}
& + \sum_{i=1}^{i^*(K)-1} x_{i,K}^n u_{i,K}^n \overline{\Delta x_i^n} + x_{i^*(K),K}^n u_{i^*(K),K}^n \left[\frac{\Delta x_{i^*(K)-1}^n}{2} + \frac{\Delta x_{i^*(K)}^n}{2} \right] \\
& - \int_0^{s(t')} \xi u(\xi, t') d\xi \Big| = \left| x_{0,K}^n u_{0,K}^n \frac{\Delta x_0^n}{2} + \sum_{i=1}^{i^*(K)-1} x_{i,K}^n u_{i,K}^n \overline{\Delta x_i^n} \right. \\
& \quad \left. + x_{i^*(K),K}^n u_{i^*(K),K}^n \left[\frac{\Delta x_{i^*(K)-1}^n}{2} + \frac{\Delta x_{i^*(K)}^n}{2} \right] \right. \\
& \quad \left. + x_{i^*(K),K}^n u_{i^*(K),K}^n [s(t') - x_{i^*(K)}^n] - x_{i^*(K),K}^n u_{i^*(K),K}^n [s(t') \right. \\
& \quad \left. - x_{i^*(K)}^n] \right| \leq \left| x_{0,K}^n u_{0,K}^n \frac{\Delta x_0^n}{2} + \sum_{i=1}^{i^*(K)-1} x_{i,K}^n u_{i,K}^n \overline{\Delta x_i^n} \right. \\
& \quad \left. + x_{i^*(K),K}^n u_{i^*(K),K}^n \left[\frac{\Delta x_{i^*(K)-1}^n}{2} + s(t') - x_{i^*(K)}^n \right] \right. \\
& \quad \left. - \int_0^{s(t')} \xi u(\xi, t') d\xi \right| + \left| x_{i^*(K),K}^n u_{i^*(K),K}^n \left[\frac{\Delta x_{i^*(K)}^n}{2} - s(t') \right. \right. \\
& \quad \left. \left. + x_{i^*(K)}^n \right] \right| \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.
\end{aligned}$$

Thus $|I[u^n(K')] - \int_0^{s(t')} \xi u(\xi, t') d\xi| \rightarrow 0$ as $n \rightarrow \infty$.

Note that in going from the third inequality to the fourth we have used that the first term in the third inequality is the difference between an integral and its Riemann sum. Thus for n sufficiently large this term can be made less than $\frac{\varepsilon}{2}$. Further the remaining term is also less than $\frac{\varepsilon}{2}$ since $x_{i^*(K),K}^n u_{i^*(K),K}^n$ is bounded and $[\frac{\Delta x_{i^*(K)}^n}{2}$

$- s(t') + x_{i^*(K)}^n] \rightarrow 0$ as $n \rightarrow \infty$.

One can now show that $I[\tilde{U}^n(K)]$ converges to $\int_0^{s(t')} \xi u(\xi, t') d\xi$ as $n \rightarrow \infty$. First observe that

$$\begin{aligned}
 (4.19) \quad & |I[\tilde{U}^n(K)] - \int_0^{s(t')} \xi u(\xi, t') d\xi| = |x_0^n (\tilde{U}_{0,K}^n - u_{0,K}^n) \frac{\Delta x_0^n}{2} \\
 & + \sum_{i=1}^{M+L(K)-1} x_i^n (\tilde{U}_{i,K}^n - u_{i,K}^n) \overline{\Delta x_i^n} \\
 & + x_{M+L(K)}^n (\tilde{U}_{M+L(K),K}^n - u_{M+L(K),K}^n) \frac{\Delta x_{M+L(K)-1}^n}{2} \\
 & + |I[u^n(K)] - I[u^n(K')]| + |I[u^n(K')] - \int_0^{s(t')} \xi u(\xi, t') d\xi| \\
 & \leq x_0^n |\tilde{U}_{0,K}^n - u_{0,K}^n| \frac{\Delta x_0^n}{2} + \sum_{i=1}^{M+L(K)-1} x_i^n |\tilde{U}_{i,K}^n - u_{i,K}^n| \overline{\Delta x_i^n} \\
 & + x_{M+L(K)}^n |\tilde{U}_{M+L(K),K}^n - u_{M+L(K),K}^n| \frac{\Delta x_{M+L(K)-1}^n}{2} \\
 & + |I[u^n(K)] - I[u^n(K')]| + |I[u^n(K')] - \int_0^{s(t')} \xi u(\xi, t') d\xi|.
 \end{aligned}$$

Recall that $\tilde{U}_{i,j}^n$ converges uniformly to u . Using this fact and

proceeding much as we did in treating the term $\sum_{i \in N_1(K)} x_i^n |u_{i,K}^n$

$- u_{i,K'}^n | \overline{\Delta x_i^n}$ in (4.18), it is now easy to see that

$$x_0^n |\tilde{U}_{0,K}^n - u_{0,K}^n| \frac{\Delta x_0^n}{2} + \sum_{i=1}^{M+L(K)-1} x_i^n |\tilde{U}_{i,K}^n - u_{i,K}^n| \overline{\Delta x_i^n}$$

$$+ x_{M+L(K)}^n |\tilde{u}_{M+L(K),K}^n - u_{M+L(K),K}^n|^{\frac{\Delta x_{M+L(K)}^n - 1}{2}} \rightarrow 0$$

as $n \rightarrow \infty$. (Here we have replaced the use of uniform continuity employed before by uniform convergence.) Also recall that

$$|I[u^n(K)] - I[u^n(K')]| \rightarrow 0 \text{ as } n \rightarrow \infty \text{ and } |I[u^n(K')] - \int_0^{s(t')} \xi u(\xi, t') d\xi| \rightarrow 0$$

as $n \rightarrow \infty$. Using all this in (4.19), we finally have that $I(\tilde{U}_{i,K}^n)$ converges to $\int_0^{s(t')} \xi u(\xi, t') d\xi$ as $n \rightarrow \infty$, as claimed.

Now we return to (4.14). First we prove that S_K^n converges to $s(t')$ as $n \rightarrow \infty$. Since $s^n(t_K^n) = S_K^n$, s^n converges uniformly to s as $n \rightarrow \infty$, s is continuous, and t_K^n approaches t' as $n \rightarrow \infty$, we can write

$$\begin{aligned} |S_K^n - s(t')| &= |s^n(t_K^n) - s(t')| \\ &= |s^n(t_K^n) - s(t_K^n) + s(t_K^n) - s(t')| \\ &\leq |s^n(t_K^n) - s(t_K^n)| + |s(t_K^n) - s(t')| \\ &\leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon \end{aligned}$$

for n sufficiently large. Hence S_K^n converges to $s(t')$ as $n \rightarrow \infty$.

Next we see that $\sum_{j=1}^K f(t_j^n) \Delta t^n$ converges to $\int_0^{t'} f(\tau) d\tau$ as $n \rightarrow \infty$.

To see this, observe that

$$\begin{aligned} \sum_{j=1}^K f(t_j^n) \Delta t^n &= \sum_{j=1}^{K-1} f(t_j^n) \Delta t^n + f(t')(t' - t_{K-1}^n) \\ &\quad - f(t')(t' - t_{K-1}^n) + f(t_K^n) \Delta t^n. \end{aligned}$$

In this expression $\sum_{j=1}^{K-1} f(t_j^n) \Delta t^n + f(t')(t' - t_{K-1}^n)$ is a Riemann sum for $\int_0^{t'} f(\tau) d\tau$ and will therefore converge to the integral as $n \rightarrow \infty$.

Further it is easily seen that $[f(t_K^n) \Delta t^n - f(t')(t' - t_{K-1}^n)] \rightarrow 0$ as $n \rightarrow \infty$. Hence $\sum_{j=1}^K f(t_j^n) \Delta t^n$ converges to $\int_0^{t'} f(\tau) d\tau$ as $n \rightarrow \infty$.

Lastly, note that since

$$x_0^n \phi(x_0^n) \frac{\Delta x_0^n}{2} + \sum_{i=1}^{M-1} \{x_i^n \phi(x_i^n)\} \Delta x_i^n + x_M^n \phi(x_M^n) \frac{\Delta x_{M-1}^n}{2}$$

is a Riemann sum for $\int_0^b \xi \phi(\xi) d\xi$, it will converge to $\int_0^b \xi \phi(\xi) d\xi$ as $n \rightarrow \infty$.

Now taking the limit in (4.14) as $n \rightarrow \infty$ and using the results obtained above we get

$$\frac{1}{2}[(s(t'))^2 - b^2] - \int_0^{t'} f(\tau) d\tau - \int_0^b \xi \phi(\xi) d\xi + \int_0^{s(t')} \xi u(\xi, t') d\xi = 0$$

for $0 < t' \leq T$. Therefore the pair (s, u) satisfies (1.4). Further

by Lemma 4.3, s is Lipschitz continuous. Also u is a solution of (1.1) by hypothesis. Hence by the discussion preceding (1.4), (s,u) is a solution of (1.1), (1.2). This (s,u) must be the only solution of (1.1), (1.2) since the solution of (1.1), (1.2) is unique. Thus the proof of Theorem 4.2 is complete.

Combining Theorem 4.1 and Theorem 4.2 we obtain the following result:

Theorem 4.3. Let $(S_j^n, \tilde{U}_{i,j}^n)$ be the solution of (2.1) - (2.3) for the mesh $(\Delta x^n, \Delta t^n)$ corresponding to the appropriate element of the subsequence $\{s^n\}_{n=1}^\infty$, and let (s,u) be the solution of (1.1), (1.2).

Then $(S_j^n, \tilde{U}_{i,j}^n)$ converges to (s,u) as $n \rightarrow \infty$.

From a practical standpoint the above result is not useful since we do not know how to pick the subsequence $\{s^n\}_{n=1}^\infty$. However, it is now possible to show that the complete sequence $\{s^n\}_{n=1}^\infty$ converges and therefore the entire sequence $\{(S_j^n, \tilde{U}_{i,j}^n)\}_{n=1}^\infty$ also converges. First we show that the entire sequence $\{s^n\}_{n=1}^\infty$ converges. Suppose the whole sequence $\{s^n\}_{n=1}^\infty$ does not converge. Then it is possible to obtain two subsequences of $\{s^n\}_{n=1}^\infty$ that converge to different values at some point. Since (cf., Lemma 4.1) each subsequence would then be equicontinuous, the result of problem 18 [15, page 156] enables us to use the procedure described in this chapter to obtain two different solutions of (1.1), (1.2), which is a contradiction. Therefore, the entire sequence $\{s^n\}_{n=1}^\infty$ must converge.

Now the whole sequence $\{(S_j^n, U_{i,j}^n)\}_{n=1}^\infty$ can be shown to converge to the exact solution (s,u) by retracing our way through the proofs of Theorem 4.1 and Theorem 4.2, only this time using the entire sequence $\{s^n\}_{n=1}^\infty$. This gives us the following theorem:

Theorem 4.4. Let $\{(\Delta x^n, \Delta t^n)\}_{n=1}^\infty$ be a sequence of ordered pairs where $\{\Delta x^n\}_{n=1}^\infty$ and $\{\Delta t^n\}_{n=1}^\infty$ are nonincreasing sequences of real numbers both of which approach zero as $n \rightarrow \infty$. Let $(S_j^n, U_{i,j}^n)$ be the solution of (2.1) - (2.3) for the mesh $(\Delta x^n, \Delta t^n)$, and let (s,u) be the solution of (1.1), (1.2). Then $(S_j^n, U_{i,j}^n)$ converges to (s,u) as $n \rightarrow \infty$.

We end this chapter with a result on stability in which we derive an a priori estimate of the dependence of the points S_j on the initial and boundary data.

Lemma 4.5. Let (s_1, u_1) and (s_2, u_2) be solutions of (1.1), (1.2) corresponding to the data $\{f_1, \phi_1, b_1\}$ and $\{f_2, \phi_2, b_2\}$ satisfying the assumptions made in Chapter I and such that $b_1 \leq b_2$. Then there exists a positive constant $C = C(b, A, T)$ such that

$$\begin{aligned}
 |s_2(t) - s_1(t)| &\leq C\{(b_2 - b_1) + \int_0^t |f_2(\tau) - f_1(\tau)| d\tau \\
 (4.4) \quad &+ \int_0^{b_1} \xi |\phi_2(\xi) - \phi_1(\xi)| d\xi + \int_{b_1}^{b_2} \xi \phi_2(\xi) d\xi\}
 \end{aligned}$$

for $0 \leq t \leq T$.

Proof. See Cannon and Hill [1, Theorem 3, page 8].

Theorem 4.5. Let ε be any arbitrary positive number. Let $\{(\Delta x^n, \Delta t^n)\}_{n=1}^{\infty}$ be a sequence of ordered pairs where $\{\Delta x^n\}_{n=1}^{\infty}$ and $\{\Delta t^n\}_{n=1}^{\infty}$ are nonincreasing sequences of real numbers both of which approach zero as $n \rightarrow \infty$. Let $(S_j^n, U_{i,j}^n)$ and $(\hat{S}_j^n, \hat{U}_{i,j}^n)$ be solutions of (2.1) - (2.3) corresponding to the mesh $(\Delta x^n, \Delta t^n)$ and the data $\{f, \phi, b\}$ and $\{\hat{f}, \hat{\phi}, \hat{b}\}$ satisfying the assumptions made in Chapter I and such that $b \leq \hat{b}$. Then there exists a positive constant $C = C(b, A, T)$ such that

$$\begin{aligned} |\hat{S}_j^n - S_j^n| \leq & \varepsilon + C\{(\hat{b} - b) + \int_0^{t_j^n} |\hat{f}(\tau) - f(\tau)| d\tau \\ & + \int_0^b \xi |\hat{\phi}(\xi) - \phi(\xi)| d\xi + \int_0^{\hat{b}} \xi \hat{\phi}(\xi) d\xi\} \end{aligned}$$

for n sufficiently large and $0 \leq t_j^n \leq T$.

Proof. We have

$$\begin{aligned} |\hat{S}_j^n - S_j^n| & \leq |S_j^n - \hat{s}(t_j^n)| + |\hat{s}(t_j^n) - s(t_j^n)| \\ (4.20) \quad & + |s(t_j^n) - S_j^n| \leq \varepsilon + C\{(\hat{b} - b) + \int_0^{t_j^n} |\hat{f}(\tau) - f(\tau)| d\tau \\ & + \int_0^b \xi |\hat{\phi}(\xi) - \phi(\xi)| d\xi + \int_0^{\hat{b}} \xi \hat{\phi}(\xi) d\xi\} \end{aligned}$$

for n sufficiently large and $0 \leq t_j^n \leq T$, with $C = C(b, A, T)$ a positive constant. In (4.20) we have used Theorem 4.4 to get

$$|\hat{S}_j^n - \hat{s}(t_j^n)| + |s(t_j^n) - S_j^n| \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

for n sufficiently large and $0 \leq t_j^n \leq T$. Also we have used Lemma 4.5 to obtain

$$\begin{aligned} |\hat{s}(t_j^n) - s(t_j^n)| &\leq C\{(\hat{b} - b) + \int_0^{t_j^n} |\hat{f}(\tau) - f(\tau)| d\tau \\ &\quad + \int_0^b \xi |\hat{\phi}(\xi) - \phi(\xi)| d\xi + \int_0^{\hat{b}} \xi \hat{\phi}(\xi) d\xi\} \end{aligned}$$

for $0 \leq t_j^n \leq T$.

CHAPTER V

A NUMERICAL EXAMPLE

In this chapter we use the numerical scheme (2.1) - (2.3) found in Chapter II to obtain a numerical solution of the following Stefan problem:

$$(5.1) \quad \left\{ \begin{array}{l} u_{xx} - u_t = 0, \quad 0 < x < s(t), \quad 0 < t \leq 2.5, \\ u(0,t) = \begin{cases} \cos(\pi/4)t, & 0 \leq t \leq 2, \\ 0, & 2 \leq t \leq 2.5, \end{cases} \\ u(s(t),t) = 0, \quad 0 \leq t \leq 2.5, \\ u(x,0) = 1 - x, \quad 0 \leq x \leq 1, \\ s(0) = 1, \end{array} \right.$$

$$(5.2) \quad \frac{ds(t)}{dt} = -u_x(s(t),t), \quad 0 < t \leq 2.5.$$

A comparison of some of the values for $U_{i,j}$ and S_j for three different mesh sizes can be found in Tables 5.1 and 5.2.

We note in Table 5.1 that for fixed (x_i, t_j) the values of $U_{i,j}$ seem to converge to some finite value as the mesh gets smaller. Also we see from Table 5.2 that for fixed t_j the values of S_j seem to converge to some finite value as the mesh approaches zero.

$\Delta x, \Delta t \backslash (x_i, t_j)$	$(.5, .25)$	$(.25, .625)$	$(.75, 1)$	$(1, 1.5)$	$(.75, 2.125)$
$\frac{1}{4}, \frac{1}{8}$	.539760	.726727	.396711	.239768	.103801
$\frac{1}{8}, \frac{1}{16}$	.537662	.727510	.398154	.242136	.102973
$\frac{1}{16}, \frac{1}{32}$	.536334	.727823	.398632	.242930	.101993

Table 5.1 Comparison of Values for $U_{i,j}$.

$\Delta x, \Delta t \backslash t_j$	.25	.875	1.25	1.875	2.125
$\frac{1}{4}, \frac{1}{8}$	1.211621	1.554632	1.708420	1.889376	1.934682
$\frac{1}{8}, \frac{1}{16}$	1.198348	1.541803	1.696362	1.877972	1.923046
$\frac{1}{16}, \frac{1}{32}$	1.192156	1.535455	1.690288	1.871998	1.916819

Table 5.2 Comparison of Values for S_j .

Let $S_j^{(\Delta x, \Delta t)}$ denote the values of S_j obtained from (2.1) - (2.3) by using the mesh $(\Delta x, \Delta t)$. Then we obtain the following estimate

$$\max_j \left| S_j^{(\frac{1}{16}, \frac{1}{32})} - S_j^{(\frac{1}{8}, \frac{1}{16})} \right| \leq 7 \times 10^{-3}.$$

LIST OF REFERENCES

- [1] J. R. CANNON and C. DENSON HILL, Existence, uniqueness, stability, and monotone dependence in a Stefan problem for the heat equation, J. Math. and Mech., 17, 1 (1967), 1-19.
- [2] J. CRANK, Two methods for the numerical solution of moving boundary problems in diffusion and heat flow, Quart. J. Mech. Appl. Math., X, 2 (1957), 220-231.
- [3] JIM DOUGLAS, JR., and T. M. GALLIE, JR., On the numerical integration of a parabolic differential equation subject to a moving boundary condition, Duke Math. J., 22 (1955), 557-572.
- [4] AVNER FRIEDMAN, Partial Differential Equations of Parabolic Type, Prentice-Hall, Englewood Cliffs, N. J., 1964, 215-237.
- [5] AVNER FRIEDMAN, Free boundary problems for parabolic equations I melting of solids, J. Math. and Mech., 8 (1959), 499-518.
- [6] AVNER FRIEDMAN, The Stefan problem in several space variables, Trans. Amer. Math. Soc., 132 (1968), 51-87.
- [7] M. GEVREY, Sur les equations aux derivees partielles du type parabolique, J. Math. Pures Appl., 9 (1913), 305-475.
- [8] C. DENSON HILL and DANIEL B. KOTLOW, Classical solutions in the large of a two-phase free boundary problem I, Arch. Rational Mech. Anal. (Berlin), 45, 1 (1972), 63-78.
- [9] C. DENSON HILL and DANIEL B. KOTLOW, Classical solutions in the large of a two-phase free boundary problem II, Arch. Rational Mech. Anal. (Berlin), 47, 5 (1972), 369-379.

- [10] HIDEO KAWARADA and MAKOTO NATORI, On numerical solutions of Stefan problem I.
- [11] GUNTER H. MEYER, Multidimensional Stefan problems, SIAM J. Numer. Anal., 10, 3 (1973), 522-536.
- [12] TATSUO NOGI, A difference scheme for solving the Stefan problem, Publ. RIMS, Kyoto Univ., 9 (1974), 543-575.
- [13] ROBERT D. RICHTMEYER and K. W. MORTON, Difference Methods for Initial Value Problems, Wiley, 1967.
- [14] MILTON E. ROSE, A method for calculating solutions of parabolic equations with a free boundary.
- [15] WALTER RUDIN, Principles of Mathematical Analysis, McGraw-Hill, New York, 1964.
- [16] WILLIAM F. TRENCH, On an explicit method for the solution of a Stefan problem, J. Soc. Indust. Appl. Math., 7, 2 (1959), 184-204.
- [17] F. P. VASILEV, On finite difference methods for the solution of Stefan's single-phase problem, J. Comp. Math. and Math. Phys., 3 (1963), 861-873.