



TEXAS TECH UNIVERSITY
Libraries™

THE NUMBER OF F_q -POINTS ON DIAGONAL HYPERSURFACES WITH MONOMIAL DEFORMATION

The Texas Tech community has made this publication openly available. [Please share](#) how this access benefits you. Your story matter to us.

Citation	Mccarthy, D.. 2024. THE NUMBER OF F_q -POINTS ON DIAGONAL HYPERSURFACES WITH MONOMIAL DEFORMATION. <i>Pacific Journal of Mathematics</i> , 328(2). https://doi.org/10.2140/pjm.2024.328.339
Citable Link	https://hdl.handle.net/2346/98677
Terms of Use	© 2024 The Author, under license to MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via Subscribe to Open. cc-by

Title page template design credit to [Harvard DASH](#).

*Pacific
Journal of
Mathematics*

**THE NUMBER OF \mathbb{F}_q -POINTS ON DIAGONAL
HYPERSURFACES WITH MONOMIAL DEFORMATION**

DERMOT MCCARTHY

THE NUMBER OF \mathbb{F}_q -POINTS ON DIAGONAL HYPERSURFACES WITH MONOMIAL DEFORMATION

DERMOT MCCARTHY

We consider the family of diagonal hypersurfaces with monomial deformation

$$D_{d,\lambda,h} : x_1^d + x_2^d + \cdots + x_n^d - d\lambda x_1^{h_1} x_2^{h_2} \cdots x_n^{h_n} = 0,$$

where $d = h_1 + h_2 + \cdots + h_n$ with $\gcd(h_1, h_2, \dots, h_n) = 1$. We first provide a formula for the number of \mathbb{F}_q -points on $D_{d,\lambda,h}$ in terms of Gauss and Jacobi sums. This generalizes a result of Koblitz, which holds in the special case $d \mid q - 1$. We then express the number of \mathbb{F}_q -points on $D_{d,\lambda,h}$ in terms of a p -adic hypergeometric function previously defined by the author. The parameters in this hypergeometric function mirror exactly those described by Koblitz when drawing an analogy between his result and classical hypergeometric functions. This generalizes a result by Sulakashna and Barman, which holds in the case $\gcd(d, q - 1) = 1$. In the special case $h_1 = h_2 = \cdots = h_n = 1$ and $d = n$, i.e., the Dwork hypersurface, we also generalize a previous result of the author which holds when q is prime.

1. Introduction

Counting the number of solutions to equations over finite fields using character sums dates back to the works of Gauss and Jacobi. A renewed interest in such problems followed subsequent important contributions from Hardy and Littlewood [1922] and Davenport and Hasse [1935]. In a seminal paper, Weil [1949] gives an exposition on the topic up to that point (as well as going on to make his famous conjectures on the zeta functions of algebraic varieties). Specifically, he develops a formula for the number of solutions over \mathbb{F}_q , the finite field with q elements, and its extensions, of $a_0x_0^{n_0} + a_1x_1^{n_1} + \cdots + a_kx_k^{n_k} = 0$, in terms of what we now call Gauss sums and Jacobi sums. The techniques involved have since become standard practice and can be found in many well-known text books, e.g., [Berndt et al. 1998; Ireland and Rosen 1990]. Since then, many authors have used and adapted the techniques outlined in Weil's paper to study other equations, e.g.,

MSC2020: primary 11G25, 33E50; secondary 11S80, 11T24, 33C99.

Keywords: diagonal hypersurface, Gauss sum, Jacobi sum, finite field hypergeometric function, p -adic hypergeometric function, counting points.

[Delsarte 1951; Furtado Gomide 1949; Koblitz 1983]. Of particular interest is the work of Koblitz [1983] where he examines the family of diagonal hypersurfaces with monomial deformation

$$(1-1) \quad D_{d,\lambda,h} : x_1^d + x_2^d + \cdots + x_n^d - d\lambda x_1^{h_1} x_2^{h_2} \cdots x_n^{h_n} = 0,$$

where $h_i \in \mathbb{Z}^+$, with $\gcd(h_1, h_2, \dots, h_n) = 1$, and $d = h_1 + h_2 + \cdots + h_n$. Koblitz's main result [1983, Theorem 2] gives a formula for the number of \mathbb{F}_q -points on $D_{d,\lambda,h}$ in the terms of Gauss and Jacobi sums, in the case $d \mid q - 1$. Using the analogy between Gauss sums and the gamma function, he notes that the main term in his formula can be considered a finite field analogue of a classical hypergeometric function. The purpose of this paper is to study $D_{d,\lambda,h}$ more generally, i.e., when the condition $d \mid q - 1$ is removed. Firstly, we generalize Koblitz's result and provide a formula for the number of \mathbb{F}_q -points on $D_{d,\lambda,h}$ in terms of Gauss and Jacobi sums without the condition $d \mid q - 1$. We then express the number of \mathbb{F}_q -points on $D_{d,\lambda,h}$ in terms of a p -adic hypergeometric function previously defined by the author. The parameters in this hypergeometric function mirror exactly those described by Koblitz when drawing an analogy between his result and classical hypergeometric functions. This generalizes a result of [Sulakashna and Barman 2022], which holds in the case $\gcd(d, q - 1) = 1$. We also examine the special case when $h_1 = h_2 = \cdots = h_n = 1$ and $d = n$, i.e., the Dwork hypersurface, and generalize a previous result of the author, which holds when q is prime.

2. Statement of results

Let $q = p^r$ be a prime power and let \mathbb{F}_q denote the finite field with q elements. Let $\widehat{\mathbb{F}_q^*}$ denote the group of multiplicative characters of \mathbb{F}_q^* . We extend the domain of $\chi \in \widehat{\mathbb{F}_q^*}$ to \mathbb{F}_q by defining $\chi(0) := 0$ (including for the trivial character ε) and denote $\bar{\chi}$ as the inverse of χ . Let T be a fixed generator of $\widehat{\mathbb{F}_q^*}$. Let θ be a fixed nontrivial additive character of \mathbb{F}_q and for $\chi \in \widehat{\mathbb{F}_q^*}$ we define the Gauss sum $g(\chi) := \sum_{x \in \mathbb{F}_q} \chi(x)\theta(x)$. For $\chi_1, \chi_2, \dots, \chi_k \in \widehat{\mathbb{F}_q^*}$, we define the Jacobi sum $J(\chi_1, \chi_2, \dots, \chi_k) := \sum_{t_i \in \mathbb{F}_q, t_1 + t_2 + \cdots + t_k = 1} \chi_1(t_1)\chi_2(t_2) \cdots \chi_k(t_k)$.

We consider the family of diagonal hypersurfaces with monomial deformation described in (1-1). Let $t := \gcd(d, q - 1)$ throughout and define

$$(2-1) \quad W := \left\{ w = (w_1, w_2, \dots, w_n) \in \mathbb{Z}^n : 0 \leq w_i < t, \sum_{i=1}^n w_i \equiv 0 \pmod{t} \right\}.$$

Define an equivalence relation \sim_h on W by

$$(2-2) \quad w \sim_h w' \text{ if } w - w' \text{ is a multiple modulo } t \text{ of } h = (h_1, h_2, \dots, h_n).$$

We denote the class containing w by $[w]$. If $h = (1, 1, \dots, 1)$ we write \sim_1 . We note, in this case, that each class contains a representative w where some $w_i = 0$,

for $1 \leq i \leq n$. We will write $[w_0]$ to indicate that we have chosen such a representative for a particular class.

Our first result provides a formula for the number of \mathbb{F}_q -points on $D_{d,\lambda,h}$ in terms of Gauss and Jacobi sums, without the condition $d \mid q - 1$. We will use $\mathbb{A}^n(\mathbb{F}_q)$ and $\mathbb{P}^n(\mathbb{F}_q)$ to denote the affine and projective n -spaces, respectively, over \mathbb{F}_q . We denote the subset of elements in these spaces where all coordinates are nonzero by $\mathbb{A}^n(\mathbb{F}_q^*)$ and $\mathbb{P}^n(\mathbb{F}_q^*)$.

Theorem 2.1. *Let $N_q(D_{d,\lambda,h})$ be the number of points in $\mathbb{P}^{n-1}(\mathbb{F}_q)$ on $D_{d,\lambda,h}$. Then*

$$N_q(D_{d,\lambda,h}) = \frac{q^{n-1} - 1}{q - 1} - \sum_{w^*} J(T^{w_1 \frac{q-1}{t}}, T^{w_2 \frac{q-1}{t}}, \dots, T^{w_n \frac{q-1}{t}}) + \frac{1}{q - 1} \sum_{s,w} \frac{g(T^{w_1 \frac{q-1}{t} + h_1 s}) g(T^{w_2 \frac{q-1}{t} + h_2 s}) \dots g(T^{w_n \frac{q-1}{t} + h_n s})}{g(T^{ds})} T^{ds}(d\lambda),$$

where the first sum is over all $w^* = (w_1, w_2, \dots, w_n) \in W$ such that $0 < w_i < t$ for all i , and the second sum is over all $s \in \{0, 1, \dots, \frac{q-1}{t} - 1\}$ and all $w = (w_1, w_2, \dots, w_n) \in W$.

Theorem 2.1 generalizes [Koblitz 1983, Theorem 2], which holds in the case $d \mid q - 1$. Using an analogy between Gauss sums and the gamma function, Koblitz noted that the second summand in his formula, which corresponds to the second summand in Theorem 2.1 above with $t = d$, can be considered a finite field analogy of the classical hypergeometric function

$$(2-3) \quad \prod_{i=1}^n \Gamma\left(\frac{w_i}{d}\right) \cdot {}_dF_{d-1} \left[\begin{matrix} \dots \dots \frac{w_i}{dh_i} + \frac{b_i}{h_i} \dots \dots \\ \frac{1}{d} \quad \frac{2}{d} \quad \dots \quad \frac{d-1}{d} \end{matrix} \middle| \lambda^d h_1^{h_1} \dots h_n^{h_n} \right],$$

where the top line parameters range over all $i = 1, \dots, n$ and, for each i , all $b_i = 0, \dots, h_i - 1$. The main purpose of this paper is to express $N_q(D_{d,\lambda,h})$ in terms of a p -adic hypergeometric function previously defined by the author, whereby the parameters in this p -adic hypergeometric function mirror exactly those described by Koblitz in (2-3) above.

Next, we rewrite Theorem 2.1 in a way more amenable to manipulation when we pass to the p -adic setting.

Corollary 2.2.

$$N_q(D_{d,\lambda,h}) = \frac{q^{n-1} - 1}{q - 1} - \frac{1}{q} \sum_{\substack{w \in W \\ \text{some } w_i = 0}} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t}}) + \frac{1}{q(q-1)} \sum_{s,w} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t} + h_i s}) g(T^{-ds}) T^{ds}(-d\lambda)$$

where the first sum is over all $w = (w_1, w_2, \dots, w_n) \in W$ such that at least one $w_i = 0$, and the second sum is over **either** all $s \in \{0, 1, \dots, \frac{q-1}{r} - 1\}$ and all $w = (w_1, w_2, \dots, w_n) \in W$ **or** all $s \in \{0, 1, \dots, q - 2\}$ and all $w = (w_1, w_2, \dots, w_n) \in W / \sim_h$. In the latter case, the sum is independent of the choice of equivalence class representatives.

We now define our p -adic hypergeometric function. Let \mathbb{Z}_p denote the ring of p -adic integers, \mathbb{Q}_p the field of p -adic numbers, $\overline{\mathbb{Q}_p}$ the algebraic closure of \mathbb{Q}_p , and \mathbb{C}_p the completion of $\overline{\mathbb{Q}_p}$. Let \mathbb{Z}_q be the ring of integers in the unique unramified extension of \mathbb{Q}_p with residue field \mathbb{F}_q . Recall that for each $x \in \mathbb{F}_q^*$, there is a unique Teichmüller representative $\omega(x) \in \mathbb{Z}_q^\times$ such that $\omega(x)$ is a $(q-1)$ -st root of unity and $\omega(x) \equiv x \pmod{p}$. Therefore, we define the Teichmüller character to be the primitive character $\omega : \mathbb{F}_q^* \rightarrow \mathbb{Z}_q^\times$ given by $x \mapsto \omega(x)$, which we extend with $\omega(0) := 0$.

Definition 2.3 [McCarthy 2013, Definition 5.1]. Let $q = p^r$ for p an odd prime. Let $\lambda \in \mathbb{F}_q$, $m \in \mathbb{Z}^+$ and $a_i, b_i \in \mathbb{Q} \cap \mathbb{Z}_p$ for $1 \leq i \leq m$. Then define

$$\begin{aligned}
 & {}_mG_m \left[\begin{matrix} a_1, a_2, \dots, a_m \\ b_1, b_2, \dots, b_m \end{matrix} \middle| \lambda \right]_q \\
 & := \frac{-1}{q-1} \sum_{s=0}^{q-2} (-1)^{sm} \overline{\omega}^s(\lambda) \\
 & \quad \times \prod_{i=1}^m \prod_{k=0}^{r-1} \frac{\Gamma_p(\left(\left(a_i - \frac{s}{q-1}\right)p^k\right)}{\Gamma_p(\langle a_i p^k \rangle)} \frac{\Gamma_p(\left(\left(-b_i + \frac{s}{q-1}\right)p^k\right)}{\Gamma_p(\langle -b_i p^k \rangle)} (-p)^{-\lfloor \langle a_i p^k \rangle - \frac{sp^k}{q-1} \rfloor - \lfloor \langle -b_i p^k \rangle + \frac{sp^k}{q-1} \rfloor}.
 \end{aligned}$$

We note that the value of ${}_mG_m[\dots]$ depends only on the fractional part of the a_i and b_i parameters, and is invariant if we change the order of the parameters. Our main result expresses $N_q(D_{d,\lambda,h})$ in terms of this function.

Theorem 2.4. Let $q = p^r$ for p an odd prime. Then, for $p \nmid dh_1 \cdots h_n$,

$$\begin{aligned}
 N_q(D_{d,\lambda,h}) &= \frac{q^{n-1} - 1}{q-1} - \frac{(-1)^n}{q} \sum_{\substack{w \in W \\ \text{some } w_i = 0}} C(w) \\
 & \quad + \frac{(-1)^n}{q} \sum_{[w] \in W / \sim_h} C(w) {}_dG_d \left[\begin{matrix} \dots \dots \frac{w_i}{th_i} + \frac{b_i}{h_i} \dots \dots \\ 1 \quad \frac{1}{d} \quad \frac{2}{d} \quad \dots \quad \frac{d-1}{d} \end{matrix} \middle| (\lambda^d h_1^{h_1} \cdots h_n^{h_n})^{-1} \right]_q
 \end{aligned}$$

where the top line parameters in ${}_dG_d$ are the list

$$\left[\frac{w_i}{th_i} + \frac{b_i}{h_i} \mid i = 1, \dots, n; b_i = 0, 1, \dots, h_i - 1 \right],$$

and

$$(2-4) \quad C(w) := \prod_{i=1}^n \prod_{a=0}^{r-1} \Gamma_p \left(\left(\frac{w_i}{t} \right) p^a \right) (-p) \left(\left(\frac{w_i}{t} \right) p^a \right).$$

As we can see, the parameters of ${}_dG_d$ in [Theorem 2.4](#) mirror exactly those in (2-3) (when $d \mid q - 1$ and so $t = d$) up to inversion of the argument $\lambda^d h_1^{h_1} \cdots h_n^{h_n}$. This inversion is a feature of the definition of the function ${}_mG_m$. Because we are summing over W / \sim_h , we can remove this inversion while also swapping the top and bottom line parameters, which gives a more natural representation, in the opinion of the author. This can be seen more clearly later, in [Corollary 2.9](#), where we get an all integral bottom line parameters.

Corollary 2.5. *Let $q = p^r$ for p an odd prime. Then, for $p \nmid d h_1 \cdots h_n$,*

$$N_q(D_{d,\lambda,h}) = \frac{q^{n-1} - 1}{q - 1} - \frac{(-1)^n}{q} \sum_{\substack{w \in W \\ \text{some } w_i = 0}} C(w) + \frac{(-1)^n}{q} \sum_{[w] \in W / \sim_h} C(-w) {}_dG_d \left[\begin{matrix} 1 & \frac{1}{d} & \frac{2}{d} & \cdots & \frac{d-1}{d} \\ \dots & \frac{w_i}{th_i} + \frac{b_i}{h_i} & \dots & \dots & \dots \end{matrix} \middle| \lambda^d h_1^{h_1} \cdots h_n^{h_n} \right]_q.$$

Ideally, in [Theorem 2.4](#) and [Corollary 2.5](#), we would like to combine both sums into a single hypergeometric term. In general, it seems that this is not possible. However, it can be achieved in two special cases as we see in the next two results. The first is when $\gcd(d, q - 1) = 1$ and the second is when all $h_i = 1$, i.e., the Dwork hypersurface.

Corollary 2.6. *Let $q = p^r$ for p an odd prime. If $\gcd(d, q - 1) = 1$ then, for $p \nmid d h_1 \cdots h_n$,*

$$N_q(D_{d,\lambda,h}) = \frac{q^{n-1} - 1}{q - 1} + (-1)^n {}_{d-1}G_{d-1} \left[\begin{matrix} \frac{1}{d} & \frac{2}{d} & \cdots & \frac{d-1}{d} \\ \dots & \frac{b_i}{h_i} & \dots & \dots \end{matrix} \middle| \lambda^d h_1^{h_1} \cdots h_n^{h_n} \right]_q$$

where the bottom line parameters in ${}_{d-1}G_{d-1}$ are the list

$$\left[\frac{b_i}{h_i} \mid i = 1, \dots, n; b_i = 0, 1, \dots, h_i - 1 \right]$$

with exactly one zero removed.

[Corollary 2.6](#) is Theorem 1.2 of [\[Sulakashna and Barman 2022\]](#).

When $h_1 = h_2 = \cdots = h_n = 1$ and $d = n$ in (1-1), we recover the Dwork hypersurface, which we will denote D_λ , i.e.,

$$D_\lambda : x_1^n + x_2^n + \cdots + x_n^n - n\lambda x_1 x_2 \cdots x_n = 0.$$

We now provide formulas for the number of \mathbb{F}_q -points on D_λ , first in terms of Gauss and Jacobi sums, and then in terms of the p -adic hypergeometric function. For a

given $w = (w_1, w_2, \dots, w_n) \in W$, define n_k to be the number of k 's appearing in w , i.e., $n_k = |\{w_i \mid 1 \leq i \leq n, w_i = k\}|$. We then let $S_w := \{k \mid 0 \leq k \leq t - 1, n_k = 0\}$ and S_w^c denote its complement in $\{0, 1, \dots, t - 1\}$. So the elements of S_w are the numbers from 0 to $t - 1$, inclusive, which do not appear in w . We define the following lists:

$$(2-5) \quad A_w : \left[\frac{t-k}{t} \mid k \in S_w \right] \cup \left[\frac{b}{n} \mid 0 \leq b \leq n - 1, b \not\equiv 0 \pmod{\frac{n}{t}} \right];$$

$$(2-6) \quad B_w : \left[\frac{t-k}{t} \text{ repeated } n_k - 1 \text{ times} \mid k \in S_w^c \right].$$

We note both lists contain $n - |S_w^c|$ numbers.

Corollary 2.7 (corollary to [Theorem 2.1](#)). *Let $N_q(D_\lambda)$ be the number of points in $\mathbb{P}^{n-1}(\mathbb{F}_q)$ on D_λ . Let $t = \gcd(n, q - 1)$. Then, for $\lambda \neq 0$,*

$$\begin{aligned} &N_q(D_\lambda) \\ &= \frac{q^{n-1} - 1}{q - 1} + \frac{1}{q(q-1)} \sum_{s,w} \left[\prod_{k \in S_w^c} \frac{g(T^{k \frac{q-1}{t} + s})^{n_k - 1}}{g(T^{-k \frac{q-1}{t} - s})} T^{k \frac{q-1}{t} + s} (-1)^q \right] g(T^{-ns}) T^{ns} (-n\lambda) \end{aligned}$$

where the sum is over *either* all $s \in \{0, 1, \dots, \frac{q-1}{t} - 1\}$ and all $w = (w_1, w_2, \dots, w_n) \in W$ or all $s \in \{0, 1, \dots, q - 2\}$ and all $w = (w_1, w_2, \dots, w_n) \in W / \sim_1$. In the latter case, the sum is independent of the choice of equivalence class representatives.

Theorem 2.8. *Let $q = p^r$ for p an odd prime. Let $N_q(D_\lambda)$ be the number of points in $\mathbb{P}^{n-1}(\mathbb{F}_q)$ on D_λ for some $\lambda \in \mathbb{F}_q^*$. Let $t = \gcd(n, q - 1)$ and let $C(w)$ be defined by (2-4). Then, for $p \nmid n$,*

$$N_q(D_\lambda) = \frac{q^{n-1} - 1}{q - 1} + (-1)^n \sum_{[w_0] \in W / \sim_1} C(w_0) {}_tG_l \left[\begin{matrix} A_{w_0} \\ B_{w_0} \end{matrix} \mid \lambda^n \right]_q.$$

[Theorem 2.8](#) generalizes [Theorem 2.2](#) in [\[McCarthy 2017\]](#) which holds for $q = p$. Finally, if we let $\gcd(n, q - 1) = 1$ in [Theorem 2.8](#), or we let $h_1 = h_2 = \dots = h_n = 1$ in [Corollary 2.6](#), it easy to see that we arrive at the following result.

Corollary 2.9. *If $\gcd(n, q - 1) = 1$ then, for $p \nmid n$,*

$$N_q(D_\lambda) = \frac{q^{n-1} - 1}{q - 1} + (-1)^n {}_{n-1}G_{n-1} \left[\begin{matrix} \frac{1}{n} & \frac{2}{n} & \dots & \frac{n-1}{n} \\ 1 & 1 & \dots & 1 \end{matrix} \mid \lambda^n \right]_q.$$

[Corollary 2.9](#) generalizes [Corollary 2.3](#) in [\[McCarthy 2017\]](#) which holds for $q = p$.

3. Preliminaries

We start by recalling some properties of Gauss and Jacobi sums. See [\[Berndt et al. 1998; Ireland and Rosen 1990\]](#) for further details, noting that we have adjusted

results to take into account $\varepsilon(0) = 0$, where ε is the trivial character. We first note that $G(\varepsilon) = -1$. For $\chi \in \widehat{\mathbb{F}_q^*}$,

$$(3-1) \quad G(\chi)G(\bar{\chi}) = \begin{cases} \chi(-1)q & \text{if } \chi \neq \varepsilon, \\ 1 & \text{if } \chi = \varepsilon. \end{cases}$$

For $\chi_1, \chi_2, \dots, \chi_k \in \widehat{\mathbb{F}_q^*}$ and $\alpha \in \mathbb{F}_q$, we define the generalized Jacobi sum

$$J_\alpha(\chi_1, \chi_2, \dots, \chi_k) := \sum_{t_i \in \mathbb{F}_q, t_1+t_2+\dots+t_k=\alpha} \chi_1(t_1)\chi_2(t_2) \cdots \chi_k(t_k).$$

When $\alpha = 1$ we recover the usual Jacobi sum as defined in Section 2.

Proposition 3.1. For $\chi_1, \chi_2, \dots, \chi_k \in \widehat{\mathbb{F}_q^*}$,

$$J_0(\chi_1, \chi_2, \dots, \chi_k) = \begin{cases} (q-1)^k - (q-1)J(\chi_1, \chi_2, \dots, \chi_k) & \text{if } \chi_1, \chi_2, \dots, \chi_k \text{ all trivial,} \\ -(q-1)J(\chi_1, \chi_2, \dots, \chi_k) & \text{if } \chi_1\chi_2 \cdots \chi_k \text{ trivial but at least} \\ & \text{one of } \chi_1, \chi_2, \dots, \chi_k \text{ nontrivial,} \\ 0 & \text{if } \chi_1\chi_2 \cdots \chi_k \text{ nontrivial.} \end{cases}$$

Proposition 3.2. For $\chi_1\chi_2 \cdots \chi_k$ trivial but at least one of $\chi_1, \chi_2, \dots, \chi_k$ nontrivial then

$$J(\chi_1, \chi_2, \dots, \chi_k) = -\chi_k(-1)J(\chi_1, \chi_2, \dots, \chi_{k-1}).$$

Proposition 3.3. For $\chi_1, \chi_2, \dots, \chi_k$ all trivial,

$$J(\chi_1, \chi_2, \dots, \chi_k) = [(q-1)^k + (-1)^{k+1}]/q.$$

Proposition 3.4. For $\chi_1, \chi_2, \dots, \chi_k$ not all trivial,

$$J(\chi_1, \chi_2, \dots, \chi_k) = \begin{cases} \frac{G(\chi_1)G(\chi_2) \cdots G(\chi_k)}{G(\chi_1\chi_2 \cdots \chi_k)} & \text{if } \chi_1\chi_2 \cdots \chi_k \neq \varepsilon, \\ -\frac{G(\chi_1)G(\chi_2) \cdots G(\chi_k)}{q} & \text{if } \chi_1\chi_2 \cdots \chi_k = \varepsilon. \end{cases}$$

We now recall the p -adic gamma function. For further details, see [Koblitz 1980]. Let p be an odd prime. For $n \in \mathbb{Z}^+$ we define the p -adic gamma function as

$$\Gamma_p(n) := (-1)^n \prod_{\substack{0 < j < n \\ p \nmid j}} j$$

and extend it to all $x \in \mathbb{Z}_p$ by setting $\Gamma_p(0) := 1$ and $\Gamma_p(x) := \lim_{n \rightarrow x} \Gamma_p(n)$ for $x \neq 0$, where n runs through any sequence of positive integers p -adically approaching x . This limit exists, is independent of how n approaches x , and

determines a continuous function on \mathbb{Z}_p with values in \mathbb{Z}_p^* . The function satisfies the following product formula.

Theorem 3.5 [Gross and Koblitz 1979, Theorem 3.1]. *If $h \in \mathbb{Z}^+$, $p \nmid h$ and $0 \leq x < 1$ with $(q - 1)x \in \mathbb{Z}$, then*

$$(3-2) \quad \prod_{a=0}^{r-1} \prod_{b=0}^{h-1} \Gamma_p\left(\left\langle \frac{x+b}{h} p^a \right\rangle\right) = \omega(h^{(q-1)x}) \prod_{a=0}^{r-1} \Gamma_p\left(\langle xp^a \rangle\right) \prod_{b=1}^{h-1} \Gamma_p\left(\left\langle \frac{b}{h} p^a \right\rangle\right).$$

We note that in the original statement of **Theorem 3.5** in [Gross and Koblitz 1979], ω is the Teichmüller character of \mathbb{F}_p^* . However, the result above still holds as $\omega|_{\mathbb{F}_p^*}$ is the Teichmüller character of \mathbb{F}_p^* .

The Gross–Koblitz formula allows us to relate Gauss sums and the p -adic gamma function. Let $\pi \in \mathbb{C}_p$ be the fixed root of $x^{p-1} + p = 0$ that satisfies $\pi \equiv \zeta_p - 1 \pmod{(\zeta_p - 1)^2}$.

Theorem 3.6 [Gross and Koblitz 1979, Theorem 1.7]. *For $j \in \mathbb{Z}$,*

$$g(\bar{\omega}^j) = -\pi^{(p-1) \sum_{a=0}^{r-1} \left\langle \frac{jp^a}{q-1} \right\rangle} \prod_{a=0}^{r-1} \Gamma_p\left(\left\langle \frac{jp^a}{q-1} \right\rangle\right).$$

We now recall some results of [Weil 1949; Koblitz 1983; Furtado Gomide 1949]. Note that the definitions and notation used for characters and for Gauss and Jacobi sums vary among those papers and differ from what’s defined in this paper. So, we have adjusted the statement of their results accordingly. For $d \in \mathbb{Z}^+$, let D_d denote the diagonal hypersurface

$$D_d : x_1^d + x_2^d + \dots + x_n^d = 0.$$

Theorem 3.7 [Weil 1949]. *Let $N_q^A(D_d)$ be the number of points in $\mathbb{A}^n(\mathbb{F}_q)$ on D_d . Let $t := \gcd(d, q - 1)$. Then*

$$N_q^A(D_d) = q^{n-1} - (q - 1) \sum_{w^*} J(T^{w_1 \frac{q-1}{t}}, T^{w_2 \frac{q-1}{t}}, \dots, T^{w_n \frac{q-1}{t}}),$$

where the sum is over all $w^* = (w_1, w_2, \dots, w_n) \in W$ such that $0 < w_i < t$.

Using similar methods to those in [Weil 1949; Koblitz 1983, Theorem 2] it is easy to see that

Theorem 3.8. *Let $N_q^{A,*}(D_d)$ be the number of points in $\mathbb{A}^n(\mathbb{F}_q^*)$ on D_d . Let $t := \gcd(d, q - 1)$. Then*

$$N_q^{A,*}(D_d) = \sum_{w \in W} J_0(T^{w_1 \frac{q-1}{t}}, T^{w_2 \frac{q-1}{t}}, \dots, T^{w_n \frac{q-1}{t}}).$$

The next result appears in [Koblitz 1983] in the homogenous case, and in general in [Furtado Gomide 1949]. We note that [Furtado Gomide 1949] contains a minor error. A term is omitted in the determination, but is easily fixed.

Theorem 3.9 [Furtado Gomide 1949; Koblitz 1983, Theorem 1]. *Let $N_q^{A,*}$ be the number of points in $\mathbb{A}^n(\mathbb{F}_q^*)$ on*

$$\sum_{i=1}^r a_i x_1^{m_{1i}} x_2^{m_{2i}} \dots x_n^{m_{ni}} = 0$$

for some $a_i \in \mathbb{F}_q^*$, $m_{ji} \in \mathbb{Z}_{\geq 0}$, such that for a given i , m_{ji} are not all zero. Then

$$N_q^{A,*} = \frac{1}{q} [(q-1)^n + (-1)^r (q-1)^{n-r+1}] - (q-1)^{n-r+1} \sum_{\alpha} T^{-\alpha_1}(a_1) T^{-\alpha_2}(a_2) \dots T^{-\alpha_r}(a_r) J(T^{\alpha_1}, T^{\alpha_2}, \dots, T^{\alpha_r}),$$

where the sum is over all $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_r) \neq 0$ satisfying $0 \leq \alpha_i < q-1$, $\sum_{i=1}^r \alpha_i \equiv 0 \pmod{q-1}$, and $\sum_{i=1}^r m_{ji} \alpha_i \equiv 0 \pmod{q-1}$ for all $j \in \{1, 2, \dots, n\}$.

A key step in proving the main results of this paper is to adapt [Theorem 3.9](#) to $D_{d,\lambda,h}$.

Corollary 3.10. *Let $t := \gcd(d, q-1)$. For $\lambda \neq 0$,*

$$N_q^{A,*}(D_{d,\lambda,h}) = \sum_{s,w} J(T^{w_1 \frac{q-1}{t} + h_1 s}, T^{w_2 \frac{q-1}{t} + h_2 s}, \dots, T^{w_n \frac{q-1}{t} + h_n s}) T^{ds}(d\lambda)$$

where the sum is over all $s \in \{0, 1, \dots, \frac{q-1}{t} - 1\}$ and all $w = (w_1, w_2, \dots, w_n) \in W$.

[Corollary 3.10](#) generalizes [Corollary 1](#) in [Koblitz 1983], which holds in the case $d \mid q-1$.

4. Proofs

Proof of [Corollary 3.10](#). We take $r = n+1$; $a_i = 1$, for $i = 1, \dots, n$, and $a_r = -d\lambda$; $m_{ji} = d$ if $i = j$ and zero otherwise, and, $m_{jr} = h_j$, for all $j = 1, \dots, n$, in [Theorem 3.9](#). This yields

$$(4-1) \quad N_q^{A,*}(D_{d,\lambda,h}) = \frac{1}{q} [(q-1)^n + (-1)^{n+1}] - \sum_{\alpha} T^{-\alpha_{n+1}}(-d\lambda) J(T^{\alpha_1}, T^{\alpha_2}, \dots, T^{\alpha_{n+1}})$$

where the sum is over all $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{n+1}) \neq 0$ satisfying $0 \leq \alpha_i < q-1$, $\sum_{i=1}^{n+1} \alpha_i \equiv 0 \pmod{q-1}$, and $d\alpha_j + h_j \alpha_{n+1} \equiv 0 \pmod{q-1}$ for all $j = 1, 2, \dots, n$.

The condition $d\alpha_j + h_j \alpha_{n+1} \equiv 0 \pmod{q-1}$, for all $j \in \{1, 2, \dots, n\}$, implies $t = \gcd(d, q-1)$ divides $h_j \alpha_{n+1}$ for all $j \in \{1, 2, \dots, n\}$. If l^e is a prime power dividing t but not α_{n+1} , then l divides h_j for all $j \in \{1, 2, \dots, n\}$. This is a

contradiction, as $\gcd(h_1, \dots, h_n) = 1$. Therefore, l^e divides α_{n+1} , which implies t divides α_{n+1} . So $\frac{\alpha_{n+1}}{t} \in \{0, 1, \dots, \frac{q-1}{t} - 1\}$. Let

$$s \equiv -\left(\frac{d}{t}\right)^{-1} \frac{\alpha_{n+1}}{t} \pmod{\frac{q-1}{t}}$$

such that $s \in \{0, 1, \dots, \frac{q-1}{t} - 1\}$. Then s runs around $\{0, 1, \dots, \frac{q-1}{t} - 1\}$ as $\frac{\alpha_{n+1}}{t}$ does.

We now express the conditions on α in terms of s . Firstly,

$$\begin{aligned} d\alpha_j &\equiv -h_j\alpha_{n+1} \pmod{q-1} &\implies &\frac{d}{t}\alpha_j \equiv -h_j\frac{\alpha_{n+1}}{t} \pmod{\frac{q-1}{t}} \\ & &\implies &\alpha_j \equiv h_js \pmod{\frac{q-1}{t}}. \end{aligned}$$

So $\alpha_j = h_js + w_j\frac{q-1}{t}$ for $w_j \in \{0, 1, \dots, t-1\}$, for $j \in \{1, 2, \dots, n\}$. Also,

$$(4-2) \quad \frac{\alpha_{n+1}}{t} \equiv -\left(\frac{d}{t}\right)s \pmod{\frac{q-1}{t}} \implies \alpha_{n+1} \equiv -ds \pmod{q-1}.$$

Using the fact that $\sum_{i=1}^n h_i = d$, it is easy to see that

$$(4-3) \quad \sum_{j=1}^n w_j = \sum_{j=1}^n \frac{t}{q-1}(\alpha_j - h_js) = \frac{t}{q-1} \left(\sum_{j=1}^n \alpha_j - ds \right).$$

Combining (4-2) and (4-3) we get that

$$\sum_{j=1}^n w_j \equiv 0 \pmod{t} \iff \sum_{i=1}^{n+1} \alpha_i \equiv 0 \pmod{q-1}.$$

Substituting for α , (4-1) becomes

$$(4-4) \quad N_q^{A,*}(D_{d,\lambda,h}) = \frac{1}{q} [(q-1)^n + (-1)^{n+1}] - \sum_{s,w} T^{ds}(-d\lambda) J(T^{w_1\frac{q-1}{t}+h_1s}, \dots, T^{w_n\frac{q-1}{t}+h_ns}, T^{-ds}),$$

where the sum is over all $s \in \{0, 1, \dots, \frac{q-1}{t} - 1\}$ and all $w = (w_1, w_2, \dots, w_n)$, such that $0 \leq w_i < t$ and $\sum_{i=1}^n w_i \equiv 0 \pmod{t}$, and such that not all of s, w_1, w_2, \dots, w_n are zero.

As $\sum_{i=1}^n w_i\frac{q-1}{t} + h_1s - ds \equiv 0 \pmod{q-1}$, by [Proposition 3.2](#) we have

$$J(T^{w_1\frac{q-1}{t}+h_1s}, \dots, T^{w_n\frac{q-1}{t}+h_ns}, T^{-ds}) = -J(T^{w_1\frac{q-1}{t}+h_1s}, \dots, T^{w_n\frac{q-1}{t}+h_ns}) T^{-ds}(-1),$$

and by [Proposition 3.3](#) we have

$$J(\underbrace{T^0, T^0, \dots, T^0}_{n \text{ times}}) = \frac{1}{q} [(q-1)^n + (-1)^{n+1}],$$

completing the proof. □

Proof of Theorem 2.1. We follow Koblitz [1983, Theorem 2] and note that

$$(4-5) \quad N_q(D_{d,\lambda,h}) - N_q^*(D_{d,\lambda,h}) = N_q(D_{d,0,h}) - N_q^*(D_{d,0,h}).$$

We know

$$(4-6) \quad N_q(D_{d,0,h}) = \frac{N_q^A(D_d) - 1}{q - 1} = \frac{q^{n-1} - 1}{q - 1} - \sum_{w^*} J(T^{w_1 \frac{q-1}{t}}, T^{w_2 \frac{q-1}{t}}, \dots, T^{w_n \frac{q-1}{t}})$$

by Weil’s result, Theorem 3.7 above;

$$N_q^*(D_{d,\lambda,h}) = \frac{1}{q - 1} \sum_{s,w} J(T^{w_1 \frac{q-1}{t} + h_1 s}, T^{w_2 \frac{q-1}{t} + h_2 s}, \dots, T^{w_n \frac{q-1}{t} + h_n s}) T^{ds}(d\lambda)$$

when $\lambda \neq 0$, by Corollary 3.10; and

$$N_q^*(D_{d,0,h}) = N_q^*(D_d) = \frac{1}{q - 1} \sum_w J_0(T^{w_1 \frac{q-1}{t}}, T^{w_2 \frac{q-1}{t}}, \dots, T^{w_n \frac{q-1}{t}})$$

by Theorem 3.8.

Using Propositions 3.1, 3.3 and 3.4, we get that for $\lambda \neq 0$,

$$\begin{aligned} (4-7) \quad & (q-1)(N_q^*(D_{d,\lambda,h}) - N_q^*(D_{d,0,h})) \\ &= \sum_{\substack{s,w \\ s \neq 0}} J(T^{w_1 \frac{q-1}{t} + h_1 s}, T^{w_2 \frac{q-1}{t} + h_2 s}, \dots, T^{w_n \frac{q-1}{t} + h_n s}) T^{ds}(d\lambda) \\ &+ \sum_w J(T^{w_1 \frac{q-1}{t}}, T^{w_2 \frac{q-1}{t}}, \dots, T^{w_n \frac{q-1}{t}}) - \sum_w J_0(T^{w_1 \frac{q-1}{t}}, T^{w_2 \frac{q-1}{t}}, \dots, T^{w_n \frac{q-1}{t}}) \\ &= \sum_{\substack{s,w \\ s \neq 0}} J(T^{w_1 \frac{q-1}{t} + h_1 s}, T^{w_2 \frac{q-1}{t} + h_2 s}, \dots, T^{w_n \frac{q-1}{t} + h_n s}) T^{ds}(d\lambda) \\ &+ q \sum_{\substack{w \\ w \neq 0}} J(T^{w_1 \frac{q-1}{t}}, T^{w_2 \frac{q-1}{t}}, \dots, T^{w_n \frac{q-1}{t}}) + qJ(\varepsilon, \varepsilon, \dots, \varepsilon) - (q-1)^n \\ &= \sum_{\substack{s,w \\ s \neq 0}} J(T^{w_1 \frac{q-1}{t} + h_1 s}, T^{w_2 \frac{q-1}{t} + h_2 s}, \dots, T^{w_n \frac{q-1}{t} + h_n s}) T^{ds}(d\lambda) \\ &+ q \sum_{\substack{w \\ w \neq 0}} J(T^{w_1 \frac{q-1}{t}}, T^{w_2 \frac{q-1}{t}}, \dots, T^{w_n \frac{q-1}{t}}) + (-1)^{n+1} \\ &= \sum_{s,w} \frac{g(T^{w_1 \frac{q-1}{t} + h_1 s})g(T^{w_2 \frac{q-1}{t} + h_2 s}) \dots g(T^{w_n \frac{q-1}{t} + h_n s})}{g(T^{ds})} T^{ds}(d\lambda). \end{aligned}$$

Combining (4-5), (4-6) and (4-7), which trivially hold for $\lambda = 0$ also, yields the result. □

Proof of Corollary 2.2. Applying (3-1) and Proposition 3.4 to Theorem 2.1 we get that

$$\begin{aligned}
 N_q(D_{d,\lambda,h}) &= \frac{q^{n-1} - 1}{q - 1} + \frac{1}{q} \sum_{w^*} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t}}) - \frac{1}{q-1} \sum_w \prod_{i=1}^n g(T^{w_i \frac{q-1}{t}}) \\
 &\quad + \frac{1}{q(q-1)} \sum_{\substack{s,w \\ s \neq 0}} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t} + h_i s}) g(T^{-ds}) T^{ds}(-d\lambda) \\
 &= \frac{q^{n-1} - 1}{q - 1} + \frac{1}{q} \sum_{w^*} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t}}) - \frac{1}{q-1} \sum_w \prod_{i=1}^n g(T^{w_i \frac{q-1}{t}}) \left(1 - \frac{1}{q}\right) \\
 &\quad + \frac{1}{q(q-1)} \sum_{s,w} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t} + h_i s}) g(T^{-ds}) T^{ds}(-d\lambda) \\
 &= \frac{q^{n-1} - 1}{q - 1} - \frac{1}{q} \sum_{\substack{w \\ \text{some } w_i = 0}} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t}}) \\
 &\quad + \frac{1}{q(q-1)} \sum_{s,w} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t} + h_i s}) g(T^{-ds}) T^{ds}(-d\lambda),
 \end{aligned}$$

where the last sum is over all $s \in \{0, 1, \dots, \frac{q-1}{t} - 1\}$ and all $w = (w_1, w_2, \dots, w_n) \in W$, as required. To get the alternative summation limits, we note that

$$\begin{aligned}
 (4-8) \quad &\sum_{s=0}^{\frac{q-1}{t}-1} \sum_{w \in W} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t} + h_i s}) g(T^{-ds}) T^{ds}(-d\lambda) \\
 &= \sum_{s=0}^{\frac{q-1}{t}-1} \sum_{j=0}^{t-1} \sum_{[w] \in W/\sim} g(T^{(w_i + j h_i) \frac{q-1}{t} + h_i s}) g(T^{-ds}) T^{ds}(-d\lambda) \\
 &= \sum_{s=0}^{\frac{q-1}{t}-1} \sum_{j=0}^{t-1} \sum_{[w] \in W/\sim} g(T^{w_i \frac{q-1}{t} + h_i (s + j \frac{q-1}{t})}) g(T^{-ds}) T^{ds}(-d\lambda) \\
 &= \sum_{s=0}^{q-2} \sum_{[w] \in W/\sim} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t} + h_i s}) g(T^{-ds}) T^{ds}(-d\lambda).
 \end{aligned}$$

This sum is independent of the choice of equivalence class representatives $[w]$, as changing representative can be countered by a simple change of variable in s . \square

Proof of Corollary 2.7. We start from Corollary 2.2 with $h = (1, 1, \dots, 1)$ and $d = n$, and rewrite using the notation described in Section 2, i.e.,

$$(4-9) \quad N_q(D_\lambda) = \frac{q^{n-1} - 1}{q - 1} - \frac{1}{q} \sum_{\substack{w \in W \\ 0 \in S_w^c}} \prod_{k \in S_w^c} g(T^{k \frac{q-1}{t}})^{n_k} \\ + \frac{1}{q(q-1)} \sum_{s, w} \prod_{k \in S_w^c} g(T^{k \frac{q-1}{t} + s})^{n_k} g(T^{-ns}) T^{ns} (-n\lambda),$$

where $t = \gcd(n, q-1)$, and the second sum is over all $s \in \{0, 1, \dots, \frac{q-1}{t} - 1\}$ and all $w \in W$. We proceed in the same fashion as the proof of Theorem 2.2 in [McCarthy 2017]. By (3-1) it is easy to see that

$$(4-10) \quad \sum_{\substack{w \\ 0 \in S_w^c}} \prod_{k \in S_w^c} g(T^{k \frac{q-1}{t}})^{n_k} = \sum_{\substack{w \\ 0 \in S_w^c}} \left[\prod_{k \in S_w^c} \frac{g(T^{k \frac{q-1}{t}})^{n_k - 1}}{g(T^{-k \frac{q-1}{t}})} \right] \left[\prod_{k \in S_w^c \setminus \{0\}} T^{k \frac{q-1}{t}} (-1) q \right].$$

We now focus on the second sum in (4-9). If $T^{k \frac{q-1}{t} + s} = \varepsilon$ then $k \frac{q-1}{t} + s \equiv 0 \pmod{q-1}$, which can only happen if $s \equiv 0 \pmod{\frac{q-1}{t}}$, in which case $s = 0$. So, if $s \neq 0$ then $T^{k \frac{q-1}{t} + s} \neq \varepsilon$. Again using (3-1), we see that, for $\lambda \neq 0$,

$$(4-11) \quad \sum_{w \in W} \sum_{s=0}^{\frac{q-1}{t}} \prod_{k \in S_w^c} g(T^{k \frac{q-1}{t} + s})^{n_k} g(T^{-ns}) T^{ns} (-n\lambda) \\ = \sum_{w \in W} \sum_{s=1}^{\frac{q-1}{t}} \left[\prod_{k \in S_w^c} \frac{g(T^{k \frac{q-1}{t} + s})^{n_k - 1}}{g(T^{-k \frac{q-1}{t} - s})} T^{k \frac{q-1}{t} + s} (-1) q \right] g(T^{-ns}) T^{ns} (-n\lambda) \\ - \sum_{w \in W} \left[\prod_{k \in S_w^c} \frac{g(T^{k \frac{q-1}{t}})^{n_k - 1}}{g(T^{-k \frac{q-1}{t}})} \right] \left[\prod_{k \in S_w^c \setminus \{0\}} T^{k \frac{q-1}{t}} (-1) q \right] \\ = \sum_{w \in W} \sum_{s=0}^{\frac{q-1}{t}} \left[\prod_{k \in S_w^c} \frac{g(T^{k \frac{q-1}{t} + s})^{n_k - 1}}{g(T^{-k \frac{q-1}{t} - s})} T^{k \frac{q-1}{t} + s} (-1) q \right] g(T^{-ns}) T^{ns} (-n\lambda) \\ + \sum_{w \in W} \left[\prod_{k \in S_w^c} \frac{g(T^{k \frac{q-1}{t}})^{n_k - 1}}{g(T^{-k \frac{q-1}{t}})} \right] \left[\prod_{k \in S_w^c} T^{k \frac{q-1}{t}} (-1) q - \prod_{k \in S_w^c \setminus \{0\}} T^{k \frac{q-1}{t}} (-1) q \right] \\ = \sum_{w \in W} \sum_{s=0}^{\frac{q-1}{t}} \left[\prod_{k \in S_w^c} \frac{g(T^{k \frac{q-1}{t} + s})^{n_k - 1}}{g(T^{-k \frac{q-1}{t} - s})} T^{k \frac{q-1}{t} + s} (-1) q \right] g(T^{-ns}) T^{ns} (-n\lambda) \\ + (q-1) \sum_{\substack{w \\ 0 \in S_w^c}} \left[\prod_{k \in S_w^c} \frac{g(T^{k \frac{q-1}{t}})^{n_k - 1}}{g(T^{-k \frac{q-1}{t}})} \right] \left[\prod_{k \in S_w^c \setminus \{0\}} T^{k \frac{q-1}{t}} (-1) q \right].$$

Accounting for (4-10) and (4-11) in (4-9) yields

$$N_q(D_\lambda) = \frac{q^{n-1}-1}{q-1} + \frac{1}{q(q-1)} \sum_{w \in W} \sum_{s=0}^{q-1} \left[\prod_{k \in S_w^c} \frac{g(T^{k \frac{q-1}{t} + s})^{n_k-1}}{g(T^{-k \frac{q-1}{t} - s})} T^{k \frac{q-1}{t} + s} (-1)^q \right] g(T^{-ns}) T^{ns} (-n\lambda).$$

To get the alternative summation limit, proceed in the same manner as in (4-8). \square

Proof of Theorem 2.4. We start from Corollary 2.2, which we rewrite as

$$(4-12) \quad N_q(D_{d,\lambda,h}) = \frac{q^{n-1}-1}{q-1} - \frac{1}{q} \sum_{\substack{w \in W \\ \text{some } w_i = 0}} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t}}) + \frac{1}{q(q-1)} \sum_{[w] \in W/\sim} R_{[w]},$$

where

$$R_{[w]} := \sum_{s=0}^{q-2} \prod_{i=1}^n g(T^{w_i \frac{q-1}{t} + h_i s}) g(T^{-ds}) T^{ds} (-d\lambda).$$

We note $R_{[w]}$ is independent of the choice of equivalence class representative.

We now let $T = \bar{\omega}$ and apply the Gross–Koblitz formula, Theorem 3.6, to both summands in (4-12). From the first summand we get that

$$(4-13) \quad \prod_{i=1}^n g(T^{w_i \frac{q-1}{t}}) = (-1)^n (-p)^{\sum_{i=1}^n \sum_{a=0}^{r-1} \langle (\frac{w_i}{t}) p^a \rangle} \prod_{i=1}^n \prod_{a=0}^{r-1} \Gamma_p(\langle (\frac{w_i}{t}) p^a \rangle) \\ = (-1)^n C(w).$$

The second, $R_{[w]}$, yields

$$(4-14) \quad R_{[w]} = (-1)^{n+1} \sum_{s=0}^{q-2} \left[\prod_{a=0}^{r-1} \prod_{i=1}^n \Gamma_p(\langle (\frac{w_i}{t} + \frac{h_i s}{q-1}) p^a \rangle) \right] \left[\prod_{a=0}^{r-1} \Gamma_p(\langle (\frac{-ds}{q-1}) p^a \rangle) \right] \\ \times (-p)^v \bar{\omega}^{ds} (-d\lambda),$$

where

$$v = \sum_{a=0}^{r-1} \sum_{i=1}^n \langle (\frac{w_i}{t} + \frac{h_i s}{q-1}) p^a \rangle + \sum_{a=0}^{r-1} \langle (\frac{-ds}{q-1}) p^a \rangle \\ = \sum_{a=0}^{r-1} \sum_{i=1}^n (\frac{w_i}{t} + \frac{h_i s}{q-1}) p^a + \sum_{a=0}^{r-1} (\frac{-ds}{q-1}) p^a - \sum_{a=0}^{r-1} \sum_{i=1}^n \lfloor (\frac{w_i}{t} + \frac{h_i s}{q-1}) p^a \rfloor - \sum_{a=0}^{r-1} \lfloor (\frac{-ds}{q-1}) p^a \rfloor \\ = \sum_{a=0}^{r-1} \sum_{i=1}^n (\frac{w_i}{t}) p^a - \sum_{a=0}^{r-1} \sum_{i=1}^n \lfloor (\frac{w_i}{t} + \frac{h_i s}{q-1}) p^a \rfloor - \sum_{a=0}^{r-1} \lfloor (\frac{-ds}{q-1}) p^a \rfloor \in \mathbb{Z}$$

as $\sum_{i=1}^n h_i = d$ and $\sum_{i=1}^n w_i \equiv 0 \pmod{t}$.

We will now use [Theorem 3.5](#) to expand the terms involving the p -adic gamma function in [\(4-14\)](#). Let $k \in \mathbb{Z}$ such that

$$k \leq \frac{w_i}{t} + \frac{h_i s}{q-1} < k + 1.$$

Then $0 \leq x := \frac{w_i}{t} + \frac{h_i s}{q-1} - k < 1$ and $(q-1)x \in \mathbb{Z}$. So, by [Theorem 3.5](#), with $h = h_i$ and $p \nmid h_i$,

$$\begin{aligned} \prod_{a=0}^{r-1} \prod_{b=0}^{h_i-1} \Gamma_p\left(\left(\frac{w_i}{th_i} + \frac{s}{q-1} + \frac{b-k}{h_i}\right)p^a\right) \\ = \omega\left(h_i^{w_i \frac{q-1}{t} + h_i s}\right) \prod_{a=0}^{r-1} \Gamma_p\left(\left(\frac{w_i}{t} + \frac{h_i s}{q-1}\right)p^a\right) \prod_{b=1}^{h_i-1} \Gamma_p\left(\left(\frac{b}{h_i}\right)p^a\right). \end{aligned}$$

As $\{b \mid b = 0, 1, \dots, h_i - 1\} \equiv \{b - k \mid b = 0, 1, \dots, h_i - 1\} \pmod{h_i}$ we have

$$\begin{aligned} (4-15) \quad \prod_{a=0}^{r-1} \prod_{b=0}^{h_i-1} \Gamma_p\left(\left(\frac{w_i}{th_i} + \frac{s}{q-1} + \frac{b}{h_i}\right)p^a\right) \\ = \omega\left(h_i^{w_i \frac{q-1}{t} + h_i s}\right) \prod_{a=0}^{r-1} \Gamma_p\left(\left(\frac{w_i}{t} + \frac{h_i s}{q-1}\right)p^a\right) \prod_{b=1}^{h_i-1} \Gamma_p\left(\left(\frac{b}{h_i}\right)p^a\right). \end{aligned}$$

Similarly, with $k \in \mathbb{Z}$ chosen such that $0 \leq x := \frac{w_i}{t} - k < 1$, we apply [Theorem 3.5](#) to get that

$$\begin{aligned} (4-16) \quad \prod_{a=0}^{r-1} \prod_{b=0}^{h_i-1} \Gamma_p\left(\left(\frac{w_i}{th_i} + \frac{b}{h_i}\right)p^a\right) \\ = \omega\left(h_i^{w_i \frac{q-1}{t}}\right) \prod_{a=0}^{r-1} \Gamma_p\left(\left(\frac{w_i}{t}\right)p^a\right) \prod_{b=1}^{h_i-1} \Gamma_p\left(\left(\frac{b}{h_i}\right)p^a\right). \end{aligned}$$

Combining [\(4-15\)](#) and [\(4-16\)](#) we have, for $p \nmid h_i$,

$$\begin{aligned} (4-17) \quad \prod_{a=0}^{r-1} \Gamma_p\left(\left(\frac{w_i}{t} + \frac{h_i s}{q-1}\right)p^a\right) \\ = \prod_{a=0}^{r-1} \prod_{b=0}^{h_i-1} \frac{\Gamma_p\left(\left(\frac{w_i}{th_i} + \frac{s}{q-1} + \frac{b}{h_i}\right)p^a\right)}{\Gamma_p\left(\left(\frac{w_i}{th_i} + \frac{b}{h_i}\right)p^a\right)} \prod_{a=0}^{r-1} \Gamma_p\left(\left(\frac{w_i}{t}\right)p^a\right) \bar{\omega}^s(h_i^{h_i}). \end{aligned}$$

A final application of [Theorem 3.5](#), this time with $k \in \mathbb{Z}$ such that $0 \leq x := k - \frac{ds}{q-1} < 1$ and $p \nmid d$, we get, after some simplification, that

$$(4-18) \quad \prod_{a=0}^{r-1} \Gamma_p\left(\left(\frac{-ds}{q-1}\right)p^a\right) = \prod_{a=0}^{r-1} \prod_{b=0}^{d-1} \frac{\Gamma_p\left(\left(\frac{-b}{d} - \frac{s}{q-1}\right)p^a\right)}{\Gamma_p\left(\left(\frac{-b}{d}\right)p^a\right)} \bar{\omega}^s(d^{-d}).$$

Accounting for (4-17) and (4-18) in (4-14) and making the change of variable $s \rightarrow (q-1) - s$ we get that

$$(4-19) \quad R_{[w]} = (-1)^{n+1} \prod_{i=1}^n \prod_{a=0}^{r-1} \Gamma_p \left(\left\langle \left(\frac{w_i}{t} \right) p^a \right\rangle \right) \sum_{s=0}^{q-2} (-1)^{sd} \\ \times \left[\prod_{i=1}^n \prod_{b=0}^{h_i-1} \prod_{a=0}^{r-1} \frac{\Gamma_p \left(\left\langle \left(\frac{w_i}{th_i} + \frac{b}{h_i} - \frac{s}{q-1} \right) p^a \right\rangle \right)}{\Gamma_p \left(\left\langle \left(\frac{w_i}{th_i} + \frac{b}{h_i} \right) p^a \right\rangle \right)} \right] \\ \times \left[\prod_{b=0}^{d-1} \prod_{a=0}^{r-1} \frac{\Gamma_p \left(\left\langle \left(\frac{-b}{d} + \frac{s}{q-1} \right) p^a \right\rangle \right)}{\Gamma_p \left(\left\langle \left(\frac{-b}{d} \right) p^a \right\rangle \right)} \right] (-p)^y \bar{\omega}^s \left(\left[\lambda^d \prod_{i=1}^n h_i^{h_i} \right]^{-1} \right),$$

where

$$y = \sum_{a=0}^{r-1} \sum_{i=1}^n \left(\frac{w_i}{t} \right) p^a - \sum_{a=0}^{r-1} \sum_{i=1}^n \left[\left(\frac{w_i}{t} - \frac{h_i s}{q-1} \right) p^a \right] - \sum_{a=0}^{r-1} \left[\left(\frac{ds}{q-1} \right) p^a \right],$$

and we have used the fact that $\bar{\omega}(-1) = -1$. Let

$$z := - \left[\sum_{i=1}^n \sum_{a=0}^{r-1} \sum_{b=0}^{h_i-1} \left[\left\langle \left(\frac{w_i}{th_i} + \frac{b}{h_i} \right) p^a \right\rangle - \frac{sp^a}{q-1} \right] + \sum_{a=0}^{r-1} \sum_{b=0}^{d-1} \left[\left\langle \left(\frac{-b}{d} \right) p^a \right\rangle + \frac{sp^a}{q-1} \right] \right].$$

Using the fact that $\lfloor mx \rfloor = \sum_{b=0}^{m-1} \lfloor x + \frac{b}{m} \rfloor$ we get that

$$y - z = \sum_{a=0}^{r-1} \sum_{i=1}^n \left(\frac{w_i}{t} \right) p^a - \sum_{a=0}^{r-1} \sum_{i=1}^n \sum_{b=0}^{h_i-1} \left[\left(\frac{w_i}{th_i} - \frac{s}{q-1} \right) p^a + \frac{b}{h_i} \right] - \sum_{a=0}^{r-1} \sum_{b=0}^{d-1} \left[\frac{sp^a}{q-1} + \frac{b}{d} \right] \\ + \sum_{i=1}^n \sum_{a=0}^{r-1} \sum_{b=0}^{h_i-1} \left[\left\langle \left(\frac{w_i}{th_i} + \frac{b}{h_i} \right) p^a \right\rangle - \frac{sp^a}{q-1} \right] + \sum_{a=0}^{r-1} \sum_{b=0}^{d-1} \left[\left\langle \left(\frac{-b}{d} \right) p^a \right\rangle + \frac{sp^a}{q-1} \right]$$

As $\gcd(p, d) = 1$, $\{b \mid b = 0, 1, \dots, d-1\} \equiv \{bp^a \mid b = 0, 1, \dots, d-1\} \pmod{d}$ and so

$$\sum_{b=0}^{d-1} \left[\left\langle \left(\frac{-b}{d} \right) p^a \right\rangle + \frac{sp^a}{q-1} \right] = \sum_{b=0}^{d-1} \left[\left\langle \left(\frac{b}{d} \right) p^a \right\rangle + \frac{sp^a}{q-1} \right] = \sum_{b=0}^{d-1} \left[\left\langle \frac{b}{d} \right\rangle + \frac{sp^a}{q-1} \right] = \sum_{b=0}^{d-1} \left[\frac{b}{d} + \frac{sp^a}{q-1} \right].$$

Similarly, as $\gcd(p, h_i) = 1$,

$$\sum_{b=0}^{h_i-1} \left[\left\langle \left(\frac{w_i}{th_i} + \frac{b}{h_i} \right) p^a \right\rangle - \frac{sp^a}{q-1} \right] = \sum_{b=0}^{h_i-1} \left[\left\langle \left(\frac{w_i}{th_i} \right) p^a + \frac{b}{h_i} \right\rangle - \frac{sp^a}{q-1} \right] \\ = \sum_{b=0}^{h_i-1} \left[\left(\frac{w_i}{th_i} \right) p^a + \frac{b}{h_i} - \left[\left(\frac{w_i}{th_i} \right) p^a + \frac{b}{h_i} \right] - \frac{sp^a}{q-1} \right]$$

$$\begin{aligned}
&= \sum_{b=0}^{h_i-1} \left[\left(\frac{w_i}{th_i} - \frac{s}{q-1} \right) p^a + \frac{b}{h_i} \right] - \sum_{b=0}^{h_i-1} \left[\left(\frac{w_i}{th_i} \right) p^a + \frac{b}{h_i} \right] \\
&= \sum_{b=0}^{h_i-1} \left[\left(\frac{w_i}{th_i} - \frac{s}{q-1} \right) p^a + \frac{b}{h_i} \right] - \left[\left(\frac{w_i}{t} \right) p^a \right].
\end{aligned}$$

So

$$y - z = \sum_{a=0}^{r-1} \sum_{i=1}^n \left(\frac{w_i}{t} \right) p^a - \sum_{a=0}^{r-1} \sum_{i=1}^n \left[\left(\frac{w_i}{t} \right) p^a \right] = \sum_{a=0}^{r-1} \sum_{i=1}^n \left(\frac{w_i}{t} \right) p^a.$$

Thus

$$\begin{aligned}
(4-20) \quad \frac{1}{q-1} R_{[w]} &= (-1)^n C(w) \\
&\times \frac{-1}{q-1} \sum_{s=0}^{q-2} (-1)^{sd} \left[\prod_{i=1}^n \prod_{b=0}^{h_i-1} \prod_{a=0}^{r-1} \frac{\Gamma_p \left(\left(\left(\frac{w_i}{th_i} + \frac{b}{h_i} - \frac{s}{q-1} \right) p^a \right) \right)}{\Gamma_p \left(\left(\left(\frac{w_i}{th_i} + \frac{b}{h_i} \right) p^a \right) \right)} \right] \\
&\times \left[\prod_{b=0}^{d-1} \prod_{a=0}^{r-1} \frac{\Gamma_p \left(\left(\left(\frac{-b}{d} + \frac{s}{q-1} \right) p^a \right) \right)}{\Gamma_p \left(\left(\left(\frac{-b}{d} \right) p^a \right) \right)} \right] (-p)^z \bar{\omega}^s \left(\left[\lambda^d \prod_{i=1}^n h_i^{h_i} \right]^{-1} \right) \\
&= (-1)^n C(w) {}_d G_d \left[\begin{matrix} \dots \frac{w_i}{th_i} + \frac{b}{h_i} \dots \\ 1 \quad \frac{1}{d} \quad \frac{2}{d} \quad \dots \quad \frac{d-1}{d} \end{matrix} \middle| (\lambda^d h_1^{h_1} \dots h_n^{h_n})^{-1} \right]_q.
\end{aligned}$$

Substituting for (4-13) and (4-20) in (4-12), we get the required result. \square

Proof of Corollary 2.5. In Theorem 2.4, we make the change of variables $w \rightarrow -w \pmod{t}$, which is a bijection on W / \sim , and $s \rightarrow (q-1) - s$ in the expansion of ${}_d G_d$ by definition. \square

Proof of Corollary 2.6. If $t = \gcd(d, q-1) = 1$ then $w = (0, 0, \dots, 0)$ is the only element in W and $C(0) = 1$. So, by Corollary 2.5

$$N_q(D_{d,\lambda,h}) = \frac{q^{n-1} - 1}{q-1} + \frac{(-1)^n}{q} \left(-1 + {}_d G_d \left[\begin{matrix} 0 \quad \frac{1}{d} \quad \frac{2}{d} \quad \dots \quad \frac{d-1}{d} \\ \dots \dots \frac{b_i}{h_i} \dots \dots \end{matrix} \middle| \lambda^d h_1^{h_1} \dots h_n^{h_n} \right]_q \right).$$

The first bottom line parameter in ${}_d G_d$ is $\frac{0}{h_1} = 0$. We will “cancel” the zero from both top and bottom to get the required ${}_{d-1} G_{d-1}$. From Definition 2.3 we see that the contribution to the summand of the top and bottom line zero is

$$\prod_{k=0}^{r-1} \frac{\Gamma_p \left(\left(\left(0 - \frac{s}{q-1} \right) p^k \right) \right)}{\Gamma_p \left(\left(0 p^k \right) \right)} \frac{\Gamma_p \left(\left(\left(0 + \frac{s}{q-1} \right) p^k \right) \right)}{\Gamma_p \left(\left(0 p^k \right) \right)} (-p)^{- \lfloor (0 p^k) - \frac{sp^k}{q-1} \rfloor - \lfloor (0 p^k) + \frac{sp^k}{q-1} \rfloor}$$

which, by Theorem 3.6 and (3-1), equals

$$g(\bar{\omega}^{-s}) g(\bar{\omega}^s) = \begin{cases} \bar{\omega}^s (-1)^q & \text{if } s \neq 0, \\ 1 & \text{if } s = 0. \end{cases}$$

We also note that when $s = 0$ the summand in [Definition 2.3](#) equals 1. Therefore,

$${}_d G_d \left[\begin{array}{c} 0, a_2, \dots, a_n \\ 0, b_2, \dots, b_n \end{array} \middle| \lambda \right]_q = 1 + q \cdot {}_{d-1} G_{d-1} \left[\begin{array}{c} a_2, \dots, a_n \\ b_2, \dots, b_n \end{array} \middle| \lambda \right]_q$$

as required. \square

Proof of Theorem 2.8. We start from [Corollary 2.7](#) and proceed in the same fashion as the second half of the proof of [Theorem 2.2](#) in [[McCarthy 2017](#)]. We let $T = \bar{\omega}$ and apply the Gross–Koblitz formula, [Theorem 3.6](#), to get

$$(4-21) \quad N_q(D_\lambda) = \frac{q^{n-1} - 1}{q - 1} + \frac{1}{q(q-1)} \sum_{[w] \in W/\sim_1} R_{[w]},$$

where

$$\begin{aligned} R_{[w]} &= \sum_{s=0}^{q-2} (-1)^{n+1} (-p)^v \bar{\omega}^{ns} (-n\lambda) \prod_{a=0}^{r-1} \Gamma_p \left(\left(\frac{-ns}{q-1} \right) p^a \right) \\ &\quad \times \prod_{k \in S_w^c} \bar{\omega} (-1)^{k \frac{q-1}{t} + s} q \prod_{a=0}^{r-1} \frac{\Gamma_p \left(\left(\frac{k}{t} + \frac{s}{q-1} \right) p^a \right)^{n_{k-1}}}{\Gamma_p \left(\left(\frac{-k}{t} - \frac{s}{q-1} \right) p^a \right)} \end{aligned}$$

with

$$\begin{aligned} v &= \sum_{k \in S_w^c} \frac{n_k k}{t} \sum_{a=0}^{r-1} p^a - \sum_{k \in S_w^c} (n_k - 1) \sum_{a=0}^{r-1} \left[\left(\frac{k}{t} + \frac{s}{q-1} \right) p^a \right] \\ &\quad + \sum_{k \in S_w^c} \sum_{a=0}^{r-1} \left[\left(-\frac{k}{t} - \frac{s}{q-1} \right) p^a \right] - \sum_{a=0}^{r-1} \left[\left(\frac{-ns}{q-1} \right) p^a \right]. \end{aligned}$$

As $p \nmid n$ we derive from [\(4-18\)](#) that

$$\begin{aligned} &\prod_{a=0}^{r-1} \Gamma_p \left(\left(\frac{-ns}{q-1} \right) p^a \right) \\ &= \prod_{a=0}^{r-1} \frac{\prod_{k=0}^{t-1} \Gamma_p \left(\left(\frac{k}{t} - \frac{s}{q-1} \right) p^a \right) \prod_{\substack{b=0 \\ b \not\equiv 0 \pmod{t}}}^{n-1} \Gamma_p \left(\left(\frac{b}{n} - \frac{s}{q-1} \right) p^a \right)}{\prod_{b=0}^{n-1} \Gamma_p \left(\left(\frac{b}{n} \right) p^a \right)} \bar{\omega}^s (n^{-n}). \end{aligned}$$

So, after some manipulation,

$$\begin{aligned}
 (4-22) \quad R_{[w]} &= (-1)^{n+1} \sum_{s=0}^{q-2} (-p)^v \bar{\omega}^{ns} (-\lambda) \left[\prod_{k \in S_w} \prod_{a=0}^{r-1} \frac{\Gamma_p\left(\left(\frac{t-k}{t} - \frac{s}{q-1}\right)p^a\right)}{\Gamma_p\left(\frac{t-k}{t}p^a\right)} \right] \\
 &\quad \times \left[\prod_{\substack{b=0 \\ b \neq 0 \pmod{\frac{n}{t}}}}^{n-1} \prod_{a=0}^{r-1} \frac{\Gamma_p\left(\left(\frac{b}{n} - \frac{s}{q-1}\right)p^a\right)}{\Gamma_p\left(\frac{b}{n}p^a\right)} \right] \\
 &\quad \times \left[\prod_{k \in S_w^c} \prod_{a=0}^{r-1} \frac{\Gamma_p\left(\left(-\frac{t-k}{t} + \frac{s}{q-1}\right)p^a\right)^{n_k-1}}{\Gamma_p\left(-\frac{t-k}{t}p^a\right)^{n_k-1}} \right] F[w],
 \end{aligned}$$

where

$$F[w] := \left[\prod_{k \in S_w^c} \prod_{a=0}^{r-1} \Gamma_p\left(\left(\frac{-k}{t}\right)p^a\right) \right]^{-1} \left[\prod_{k \in S_w^c} \prod_{a=0}^{r-1} \Gamma_p\left(\left(\frac{k}{t}\right)p^a\right)^{n_k-1} \right] \left[\prod_{k \in S_w^c} \bar{\omega}(-1)^{k \frac{q-1}{t} + s} q \right].$$

Applying the Gross–Koblitz formula, [Theorem 3.6](#), in reverse and (3-1) we get that

$$\begin{aligned}
 &\prod_{k \in S_w^c} \prod_{a=0}^{r-1} \Gamma_p\left(\left(\frac{-k}{t}\right)p^a\right) \Gamma_p\left(\left(\frac{k}{t}\right)p^a\right) \\
 &= \prod_{k \in S_w^c} g(\bar{\omega}^{-k \frac{q-1}{t}}) g(\bar{\omega}^{k \frac{q-1}{t}}) (-p)^{-\sum_{a=0}^{r-1} \left(\left(\frac{-k}{t}\right)p^a + \left(\frac{k}{t}\right)p^a\right)} \\
 &= (-1)^{r|S_w^c \setminus \{0\}|} \prod_{k \in S_w^c} \bar{\omega}(-1)^{k \frac{q-1}{t}}.
 \end{aligned}$$

Thus

$$(4-23) \quad F[w] = (-1)^{r|S_w^c \setminus \{0\}|} q^{|S_w^c|} \bar{\omega}(-1)^{s|S_w^c|} \prod_{k \in S_w^c} \prod_{a=0}^{r-1} \Gamma_p\left(\left(\frac{k}{t}\right)p^a\right)^{n_k}.$$

If we let

$$\begin{aligned}
 -z &= \sum_{k \in S_w^c} (n_k - 1) \sum_{a=0}^{r-1} \left[\left\langle -\frac{t-k}{t} p^a \right\rangle + \frac{sp^a}{q-1} \right] \\
 &\quad + \sum_{k \in S_w} \sum_{a=0}^{r-1} \left[\left\langle \frac{t-k}{t} p^a \right\rangle - \frac{sp^a}{q-1} \right] + \sum_{\substack{b=0 \\ b \neq 0 \pmod{\frac{n}{t}}}}^{n-1} \left[\left\langle \frac{b}{n} p^a \right\rangle - \frac{sp^a}{q-1} \right],
 \end{aligned}$$

then, after a lengthy but straightforward calculation, we find that

$$(4-24) \quad v - z = -r |S_w^c \setminus \{0\}| + \sum_{i=1}^n \sum_{a=0}^{r-1} \left\langle \frac{w_i}{t} p^a \right\rangle.$$

Accounting for (4-23) and (4-24) in (4-22), and then (4-21), yields the result. \square

5. Concluding remarks

When $d \mid q - 1$ it is possible express the results of Koblitz, and those in this paper, in terms of hypergeometric functions over finite fields, as defined in [Greene 1987], or using a normalized version defined in [McCarthy 2012]. For example, see [Goodson 2017a; McCarthy 2017; Nakagawa 2021] for related results. To extend these results beyond $q \equiv 1 \pmod{d}$ it is necessary to move to the p -adic setting as we have done in this paper. Other results where the p -adic hypergeometric function, ${}_mG_m$, is used to count points on certain hypersurfaces, which are special cases of the results in this paper, can be found in [Barman et al. 2016; Goodson 2017b].

References

- [Barman et al. 2016] R. Barman, H. Rahman, and N. Saikia, “Counting points on Dwork hypersurfaces and p -adic hypergeometric functions”, *Bull. Aust. Math. Soc.* **94**:2 (2016), 208–216. [MR](#) [Zbl](#)
- [Berndt et al. 1998] B. C. Berndt, R. J. Evans, and K. S. Williams, *Gauss and Jacobi sums*, Wiley, New York, 1998. [MR](#) [Zbl](#)
- [Davenport and Hasse 1935] H. Davenport and H. Hasse, “Die Nullstellen der Kongruenzetafunktionen in gewissen zyklischen Fällen”, *J. Reine Angew. Math.* **172** (1935), 151–182. [MR](#) [Zbl](#)
- [Delsarte 1951] J. Delsarte, “Nombre de solutions des équations polynomiales sur un corps fini”, exposé 39 in *Séminaire Bourbaki*, 1948/1951, W. A. Benjamin, 1951. Reprinted as pp. 321–329 in *Séminaire Bourbaki* **1**, Soc. Math. France, Paris, 1995. [MR](#) [Zbl](#)
- [Furtado Gomide 1949] E. Furtado Gomide, “On the theorem of Artin–Weil”, *Bol. Soc. Mat. São Paulo* **4** (1949), 1–18. [MR](#) [Zbl](#)
- [Goodson 2017a] H. Goodson, “A complete hypergeometric point count formula for Dwork hypersurfaces”, *J. Number Theory* **179** (2017), 142–171. [MR](#) [Zbl](#)
- [Goodson 2017b] H. Goodson, “Hypergeometric functions and relations to Dwork hypersurfaces”, *Int. J. Number Theory* **13**:2 (2017), 439–485. [MR](#) [Zbl](#)
- [Greene 1987] J. Greene, “Hypergeometric functions over finite fields”, *Trans. Amer. Math. Soc.* **301**:1 (1987), 77–101. [MR](#) [Zbl](#)
- [Gross and Koblitz 1979] B. H. Gross and N. Koblitz, “Gauss sums and the p -adic Γ -function”, *Ann. of Math. (2)* **109**:3 (1979), 569–581. [MR](#) [Zbl](#)
- [Hardy and Littlewood 1922] G. H. Hardy and J. E. Littlewood, “Some problems of “Partitio Numerorum”, IV: the singular series in Waring’s Problem and the value of the number $G(k)$ ”, *Math. Z.* **12**:1 (1922), 161–188. [MR](#) [Zbl](#)
- [Ireland and Rosen 1990] K. Ireland and M. Rosen, *A classical introduction to modern number theory*, 2nd ed., Graduate Texts in Mathematics **84**, Springer, 1990. [MR](#) [Zbl](#)
- [Koblitz 1980] N. Koblitz, *p -adic analysis: a short course on recent work*, London Math. Soc. Lecture Note Series **46**, Cambridge University Press, 1980. [MR](#) [Zbl](#)
- [Koblitz 1983] N. Koblitz, “The number of points on certain families of hypersurfaces over finite fields”, *Compositio Math.* **48**:1 (1983), 3–23. [MR](#) [Zbl](#)
- [McCarthy 2012] D. McCarthy, “Transformations of well-poised hypergeometric functions over finite fields”, *Finite Fields Appl.* **18**:6 (2012), 1133–1147. [MR](#) [Zbl](#)
- [McCarthy 2013] D. McCarthy, “The trace of Frobenius of elliptic curves and the p -adic gamma function”, *Pacific J. Math.* **261**:1 (2013), 219–236. [MR](#) [Zbl](#)

- [McCarthy 2017] D. McCarthy, “The number of \mathbb{F}_p -points on Dwork hypersurfaces and hypergeometric functions”, *Res. Math. Sci.* **4** (2017), art. id. 4. [MR](#) [Zbl](#)
- [Nakagawa 2021] A. Nakagawa, “Artin L -functions of diagonal hypersurfaces and generalized hypergeometric functions over finite fields”, 2021. [arXiv 2111.15054](#)
- [Sulakashna and Barman 2022] Sulakashna and R. Barman, “Number of \mathbb{F}_q -points on diagonal hypersurfaces and hypergeometric function”, 2022. [arXiv 2210.11732](#)
- [Weil 1949] A. Weil, “Numbers of solutions of equations in finite fields”, *Bull. Amer. Math. Soc.* **55** (1949), 497–508. [MR](#) [Zbl](#)

Received December 9, 2023.

DERMOT MCCARTHY
DEPARTMENT OF MATHEMATICS & STATISTICS
TEXAS TECH UNIVERSITY
LUBBOCK, TX
UNITED STATES
dermot.mccarthy@ttu.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2024 is US \$645/year for the electronic version, and \$875/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 328 No. 2 February 2024

Hankel operators on $L^p(\mathbb{R}_+)$ and their p -completely bounded multipliers	193
LORIS ARNOLD, CHRISTIAN LE MERDY and SAFOURA ZADEH	
Stable functorial equivalence of blocks	217
SERGE BOUC and DENIZ YILMAZ	
Lorentz–Shimogaki and Arazy–Cwikel theorems revisited	227
LÉONARD CADILHAC, FEDOR SUKOCHEV and DMITRIY ZANIN	
Finite axiomatizability of the rank and the dimension of a pro- π group	255
MARTINA CONTE and BENJAMIN KLOPSCH	
Elliptic genus and string cobordism at dimension 24	275
FEI HAN and RUIZHI HUANG	
The domination monoid in henselian valued fields	287
MARTIN HILS and ROSARIO MENNUNI	
Inverse semigroup from metrics on doubles, III: Commutativity and (in)finiteness of idempotents	325
VLADIMIR MANUILOV	
The number of \mathbb{F}_q -points on diagonal hypersurfaces with monomial deformation	339
DERMOT MCCARTHY	
Deformation of pairs and semiregularity	361
TAKEO NISHINO	